

WEST Search History

DATE: Tuesday, August 17, 2004

Hide? Set
Name Query

DB=USPT; PLUR=YES; OP=AND

☐ L1 (portner or ginnny).xa.

☐ L2 L1 and botulin\$

DB=USPT,USOC,EPAB,JPAB,DWPI; PLUR=YES; OP=OR

☐ L3 ((neospora or neosporosis or neospor\$) not neosporin\$ near5 (avirulent or a-
virulent or non-virulent or attenuated or attenuation or attenuating or modified
or modification of mutagenesis or mutation or mutant or alter or alteration or
altered or altering))

Hit
Count

236

7

20

END OF SEARCH HISTORY

Updated
West
D111015
12/18

[Generate Collection](#)[Print](#)**Search Results** - Record(s) 1 through 20 of 20 returned.

-
- ☐ 1. [6777192](#). 21 Sep 01; 17 Aug 04. Recombinant neospora antigens and their uses. Conrad; Patricia A., et al. 435/7.1; 435/7.21 435/7.92. G01N033/53 G01N033/567 G01N033/537.
-
- ☐ 2. [6716423](#). 10 Jul 00; 06 Apr 04. Recombinant neospora antigens and their uses. Conrad; Patricia A., et al. 424/93.1; 424/184.1 424/234.1 424/93.7. A61K039/00 A61K039/38 A61K039/02 A01N063/00 A01N065/00.
-
- ☐ 3. [6656479](#). 12 Sep 01; 02 Dec 03. Attenuated live neospora vaccine. Brake; David A, et al. 424/269.1; 424/258.1 424/271.1 424/273.1 424/93.1 424/93.2 435/258.1 435/69.1. A61K039/002.
-
- ☐ 4. [6600027](#). 25 Mar 99; 29 Jul 03. Polynucleotide molecules encoding neospora proteins. Krishnan; B. Rajendra, et al. 536/23.1; 435/252.3 435/320.1 435/69.1 536/23.4 536/23.5 536/24.32. C07H021/02 C07H021/04 C12P021/06 C12N001/20 C12N015/00.
-
- ☐ 5. [6436410](#). 02 Dec 98; 20 Aug 02. DNA encoding neospora dihydrofolate reductase-thymidylate synthase. Krishnan; B. Rajendra, et al. 424/265.1; 424/184.1 424/191.1 424/200.1 424/273.1 435/6 435/91.4 536/23.1 536/23.7. A61K039/002 A61K039/012 C07H021/04 C07H021/02 C12Q001/68.
-
- ☐ 6. [6376196](#). 30 Mar 99; 23 Apr 02. Recombinant neospora antigens and their uses. Conrad; Patricia, et al. 435/7.1; 435/7.21 435/7.92. G01N033/53 G01N033/567 G01N033/537.
-
- ☐ 7. [6071737](#). 16 Mar 98; 06 Jun 00. Equine Neospora isolate and its uses. Marsh; Antoinette E., et al. 435/258.1;. C12N001/14.
-
- ☐ 8. [5889166](#). 10 May 96; 30 Mar 99. Recombinant neospora antigens and their uses. Conrad; Patricia A., et al. 536/23.1; 530/300 530/350 530/371. C07H021/02 A61K038/00 C07K001/00.
-
- ☐ 9. [5707617](#). 20 Oct 94; 13 Jan 98. Bovine neospora isolates. Conrad; Patricia A., et al. 424/93.1; 435/258.1. C12N001/10.
-
- ☐ 10. JP410167983A. 12 Nov 97. 23 Jun 98. ATTENUATED LIVE NEOSPORA VACCINE. DAVID, A BRAKE, et al. A61K039/00; A61K035/68 A61K039/002 A61K039/39 C12N001/00.
-
- ☐ 11. [EP001221485A2](#). 09 Mar 99. 10 Jul 02. Polynucleotide molecules encoding neospora proteins. BRAKE, DAVID ALAN, et al. C12N015/30; C12N015/85 C12N005/10 C07K014/44 C07K016/20 A61K039/002.
-
- ☐ 12. [EP000953641A2](#). 09 Mar 99. 03 Nov 99. Polynucleotide molecules encoding neospora proteins. BRAKE, DAVID ALAN, et al. C12N015/30; C07K014/44 C07K016/20 C12N005/16 A61K039/002.
-
- ☐ 13. [EP000924295A2](#). 26 Nov 98. 23 Jun 99. DNA encoding neospora dihydrofolate reductase-thymidylate synthase. KRISHNAN, BALAKRISHNAN RAJENDRA, et al. C12N009/06; C12N015/53 C12N015/73 C07K016/40 C12N001/11 A61K039/002.
-
- ☐ 14. [EP000841392A2](#). 14 Oct 97. 13 May 98. Attenuated live neospora vaccine. BRAKE, DAVID,
-

et al. C12N001/10; A61K039/002 A61K039/00.

- ☐ 15. US20030044431A. New overexpressing homologous antigen vaccines, useful for immunization, prophylaxis or treatment of a vertebrate at risk of or suffering from, e.g. brucellosis, paratuberculosis, tuberculosis or neosporosis. BOYLE, S M, et al. A61K039/10.
- ☐ 16. EP 1221485A. Novel *Neospora caninum* GRA2 protein useful for producing vaccines against neosporosis and as diagnostic reagents. BRAKE, D A, et al. A61K039/002 C07K014/44 C07K016/20 C12N005/10 C12N015/30 C12N015/85.
- ☐ 17. EP 1221486A. Novel *Neospora caninum* SAG1 protein useful for producing vaccines against neosporosis and as diagnostic reagents. BRAKE, D A, et al. A61K039/002 C07K014/44 C07K016/20 C12N005/10 C12N015/30 C12N015/85.
- ☐ 18. EP 953641A. Polypeptides encoding *Neospora caninum* proteins, useful for vaccines against neosporosis and as diagnostic reagents. BRAKE, D A, et al. A61K031/00 A61K031/70 A61K035/12 A61K038/00 A61K038/17 A61K039/002 A61K039/02 A61K039/39 A61K048/00 A61P033/02 C07H021/00 C07H021/02 C07H021/04 C07K014/195 C07K014/44 C07K016/18 C07K016/20 C12N000/00 C12N001/11 C12N001/20 C12N001/21 C12N005/16 C12N015/00 C12N015/02 C12N015/09 C12N015/30 C12N015/63 C12P021/02 C12P021/06 C12P021/08 C12Q001/68.
- ☐ 19. EP 924295A. New dihydrofolate reductase-thymidylate synthase (DHFR-TS) protein and gene, useful in genetic constructs for preparing a live vaccine against *Neospora*. DURTSCHI, B A, et al. A45C011/00 A45C011/36 A61K031/00 A61K035/66 A61K039/00 A61K039/002 A61K039/012 C07H021/02 C07H021/04 C07K014/44 C07K016/20 C07K016/40 C12N000/00 C12N001/11 C12N001/19 C12N001/21 C12N001/36 C12N005/10 C12N009/00 C12N009/06 C12N015/09 C12N015/30 C12N015/52 C12N015/53 C12N015/54 C12N015/60 C12N015/73 C12N015/86 C12P019/34 C12P021/02 C12Q001/68 C12R001/90 G01N033/53 G01N033/531 G01N033/569 C12N001/21 C12R001:19 C12N005/10 C12R001:90.
- ☐ 20. NZ 329095A. Live attenuated *Neospora* vaccine - for protecting animals against neosporosis. BLAGBURN, B, et al. A61K000/00 A61K035/68 A61K039/00 A61K039/002 A61K039/02 A61K039/38 A61K039/39 C12N001/00 C12N001/10 C12N001/11 C12N001/13 C12N001/18 C12N001/36 C12N005/06 C12N015/30 C12N001/10 C12R001:90 C12N001/10 C12R001:90.

Generate Collection

Print

Terms	Documents
L3 near5 (avirulent or a-virulent or non-virulent or attenuated or attenuation or attenuating or modified or modification of mutagenesis or mutation or mutant or alter or alteration or altered or altering)	20

[Prev Page](#) [Next Page](#) [Go to Doc#](#)

First Hit Fwd Refs

L4: Entry 1 of 20

File: USPT

Aug 17, 2004

US-PAT-NO: 6777192

DOCUMENT-IDENTIFIER: US 6777192 B2

TITLE: Recombinant neospora antigens and their uses

DATE-ISSUED: August 17, 2004

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Conrad; Patricia A.	Davis	CA		
Barr; Bradd C.	Davis	CA		
• Anderson; Mark L.	Davis	CA		
• Sverlow; Karen W.	Vacaville	CA		

ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE	CODE
The Regents of the University of California	Oakland	CA				02

APPL-NO: 09/ 957995 [PALM]

DATE FILED: September 21, 2001

PARENT-CASE:

This application is a continuation of U.S. Ser. No. 09/281,766, filed Mar. 30, 1999, now U.S. Pat. No. 6,376,196, which is a continuation-in-part of U.S. Ser. No. 08/645,951, filed May 10, 1996, now U.S. Pat. No. 5,889,166, which is a continuation-in-part of U.S. Ser. No. 08/327,516, filed Oct. 20, 1994, now U.S. Pat. No. 5,707,617, which is a continuation-in-part of U.S. Ser. No. 08/215,858, filed Mar. 21, 1994, now abandoned.

INT-CL: [07] G01 N 33/53, G01 N 33/567, G01 N 33/537US-CL-ISSUED: 435/7.1; 435/7.21, 435/7.92US-CL-CURRENT: 435/7.1; 435/7.21, 435/7.92FIELD-OF-SEARCH: 424/184.1, 435/7.1, 435/7.21, 435/7.92, 530/350

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

PAT-NO

ISSUE-DATE

PATENTEE-NAME

US-CL

4431739

February 1984

Riggs et al.

<input type="checkbox"/>	<u>5707617</u>	January 1998	Conrad et al.
<input type="checkbox"/>	<u>6376196</u>	April 2002	Conrad et al.

FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
WO 95/25541	September 1995	WO	
WO 97/39009	October 1997	WO	

OTHER PUBLICATIONS

Levinson et al (Medical Microbiology & Immunology p. 263), 1989.*
Dubey et al (Journal of Parasitology vol. 75(5) pp 765-771), 1989.*
Lindsay et al (J. of Helminthological Society of Washington vol. 57(1) pp 86-88), 1990.*
Barr et al., Congenital Neospora infection in calves born from cows that had previously aborted Neospora-infected fetuses: Four cases (1990-1992), JAVMA 202:1: 113-117 (1993).
Anderson, et al. J. Am. Vet Med. Assoc., 198:241-244 (1991).
Barr, et al. J. Vet. Diagn. Invest., 3:39-46 (1991).
Barr, et al. Vet. Pathol., 27:354-361 (1990).
Barr, et al. Vet. Pathol., 28:110-116 (1991).
Barta, et al., Parasitology Research, 78:689-694 (1992).
Baszler, et al., J. of Clin. Micro., 34:1423-1428 (1996).
Bjerkas, et al., Clin. Diag. Lab. Immunol., 1(2):214-221 (1994).
Bjorkman, et al., Parasite Immunology, 16:643-648 (1994).
Conrad, et al., J. Vet. Diagn. Invest., 5:572-578 (1993).
Conrad, Parasitology, 106:239-249 (1993).
Dubey, et al. J. Am. Vet Med. Assoc., 193:1259-1263 (1988).
Dubey, et al. J. Am. Vet Med. Assoc., 201(5):709-713 (1992).
Dubey, et al. J. of Parasitology, 76(5):732-734 (1990).
Dubey, et al. J. Protozool. Res., 2:40-52 (1992).
Lally, et al., Am. J. Trop. Med. Hyg. 53(2):288 (1995).
Lally, et al., Clin. Diag. Lab. Immunol., 3:275-279 (1996).
Lally, et al., Mol. and Biochem. Parasitology., 75:169-178 (1996).
Lindsay, et al., "Neospora-Induced Protozoal Abortions in Cattle", The Compendium, 15(6):882-889.
Lindsay & Dubey, Am. J. Vet. Res., 50:1981-1983 (1989).
Pare, et al., J. Vet Diagn. Invest., 7:352-359 (1995).
Shivaprasad, et al., Vet. Parasitology, 34:145-148 (1989).
Speer, et al., Infect. and Immun., 50:566-571 (1988).

ART-UNIT: 1645


PRIMARY-EXAMINER: Navarro; Mark

ATTY-AGENT-FIRM: Townsend and Townsend and Crew LLP

ABSTRACT:

The present invention provides isolated bovine Neospora cultures. Also provided are recombinant immunodominant Neospora antigens. The cultures and antigens are used to develop diagnostic assays for the detection of Neospora infections in cattle and other animals. Also provided are pharmaceutical compositions for the treatment and prevention of Neospora infections.

5 Claims, 6 Drawing figures

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
Hosted by CBR Canada Mirror sites: Australia Bolivia China Korea Switzerland Taiwan USA				
Search <input type="text" value="Swiss-Prot/TrEMBL"/>		<input type="button" value="Go"/>	<input type="button" value="Clear"/>	

Search in Swiss-Prot and TrEMBL for: neospora

Swiss-Prot Release 44.3 of 16-Aug-2004

TrEMBL Release 27.3 of 16-Aug-2004

- Number of sequences found in Swiss-Prot₍₃₎ and TrEMBL₍₃₉₎: **42**
- Note that the selected sequences can be saved to a file to be later retrieved; to do so, go to the bottom of this page.
- For more directed searches, you can use the Sequence Retrieval System SRS.

Search in Swiss-Prot: There are matches to 3 out of 156998 entries

1433 NEOCA (**Q25538**)

14-3-3 protein homolog. - Neospora caninum

GRA1 NEOCA (**P90661**)

Dense granule protein 1 precursor (NcDG1) (Antigen Nc4.1). {GENE: Name=DG1} - Neospora caninum

GRA2 NEOCA (**Q25540**)

Dense granule protein 2 (NcDG2) (Antigen Nc14.1). {GENE: Name=DG2} - Neospora caninum

Search in TrEMBL: There are matches to 39 out of 1379120 entries

O44063

Antigen N54 (Fragment) - Neospora sp

O61159

Surface protein Nc-p43 - Neospora caninum

O62617

P36 protein precursor (Surface protein P36) {GENE:Name=p36} - Neospora caninum

O76144

NTPase - Neospora caninum

O96451

Surface antigen p35 - Neospora caninum

Q6EEV5

Apical complex protein (Fragment) - Neospora caninum

Q6IMH5

Rbj-like protein (Fragment) {GENE:Name=Rjl} - Neospora caninum

Q6YDA6

Putative dense granule protein 3 - Neospora caninum

Q6YFA5

Small heat shock protein - Neospora caninum

Q6YFA6

Small heat shock protein (Fragment) - Neospora caninum

Q71G51

Alpha-tubulin {GENE:Name=TUBA} - Neospora caninum

Q75UX7

Chitinase 19-1 (Fragment) {GENE:Name=chi19-1} - Actinokineospora riparia

Q86M53

Putative surface antigen protein 1 {GENE:Name=SAG1} - Neospora caninum

Q8IT72

Microneme protein NcMIC11 precursor {GENE:Name=MIC11} - Neospora caninum

Q8MZJ9

Serine proteinase inhibitor PI-S - Neospora caninum

Q8WRS0

Microneme-associated protein {GENE:Name=MIC1} - Neospora caninum

Q8WSI9

Superoxide dismutase {GENE:Name=SOD} - Neospora caninum

Q962A6

MIC2-associated protein precursor {GENE:Name=M2AP} - Neospora caninum

Q9BKL9

P20 - Neospora caninum

Q9BKM0

Gra1 (Fragment) {GENE:Name=GRA1} - Neospora caninum

Q9GSV4

NcMIC10 precursor {GENE:Name=MIC10} - Neospora caninum

Q9GTB7

Myosin E (Fragment) - Neospora caninum

Q9GTB8

Myosin D (Fragment) - Neospora caninum

Q9GTB9

Myosin C (Fragment) - Neospora caninum

Q9GTC0

Myosin B (Fragment) - Neospora caninum

Q9GTC1

Myosin A (Fragment) - Neospora caninum

Q9GU48

GRA2 protein {GENE:Name=GRA2} - Neospora caninum

Q9NB95

SUL1 - Neospora caninum

Q9NJJ4

DNA dependent RNA polymerase beta subunit' (Fragment) {GENE:Name=rpoC1} - Neospora caninum

Q9NJJ5

DNA dependent RNA polymerase beta subunit {GENE:Name=rpoB} - Neospora caninum

Q9NJS9

SAG1 precursor {GENE:Name=SAG1} - Neospora hughesi

Q9TVP5

SRS2 surface antigen (Fragment) {GENE:Name=srs2} - Neospora caninum

[Q9TVV8](#)

P29 surface antigen (Fragment) {GENE:Name=SAG1} - Neospora caninum

[Q9U483](#)

Microneme protein Nc-P38 - Neospora caninum

[Q9U4R7](#)

SAG1-related sequence 2 {GENE:Name=SRS2} - Neospora hughesi

[Q9U7D4](#)

Subtilisin-like serine protease (Fragment) - Neospora caninum

[Q9U7D6](#)

Ycf24 protein (Fragment) {GENE:Name=ycf24} - Neospora caninum

[Q9U8J9](#)

Thrombospondin-related adhesive protein homolog {GENE:Name=MIC2} - Neospora caninum

[Q9UB12](#)

Surface antigen SAG1 {GENE:Name=SAG1} - Neospora caninum

in Swiss-Prot/TrEMBL by AC, ID, description,
gene name, organism


**Please do NOT use any boolean operators (and,
or, etc.)**

If you would like to retrieve all the Swiss-Prot/TrEMBL entries contained in this list, you can enter a file name. These entries will then be saved to a file under this name in the directory outgoing of the [ExpASy anonymous ftp server](#), from where you can download it. (Please note that this temporary file will only be kept for 1 week.)

File name: Format: ☒ Swiss-Prot ☐ Fasta

or

[ExPASy Home page](#)[Site Map](#)[Search ExpASy](#)[Contact us](#)[Swiss-Prot](#)[Hosted by CBR Canada](#)[Mirror sites:](#)[Australia](#)[Bolivia](#)[China](#)[Korea](#)[Switzerland](#)[Taiwan](#)[USA](#)

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
--	--------------------------	-------------------------------	----------------------------	----------------------------

Search for

NiceProt

View of

Swiss-

Prot:

Q25538

[Printer-friendly view](#)
[Submit update](#)
[Quick BlastP search](#)

[\[Entry info\]](#)
[\[Name and origin\]](#)
[\[References\]](#)
[\[Comments\]](#)
[\[Cross-references\]](#)
[\[Keywords\]](#)
[\[Features\]](#)
[\[Sequence\]](#)
[\[Tools\]](#)

Note: most headings are clickable, even if they don't appear as links. They link to the [user manual](#) or [other documents](#).

Entry information

Entry name	1433_NEOCA
Primary accession number	Q25538
Secondary accession numbers	None
Entered in Swiss-Prot in	Release 36, July 1998
Sequence was last modified in	Release 36, July 1998
Annotations were last modified in	Release 36, July 1998
Name and origin of the protein	
Protein name	14-3-3 protein homolog
Synonyms	None
Gene name	None
From	Neospora caninum [TaxID: 29176]
Taxonomy	Eukaryota ; Alveolata ; Apicomplexa ; Coccidia ; Eimeriida ; Sarcocystidae ; Neospora .

References

[1] SEQUENCE FROM NUCLEIC ACID.
STRAIN=Nc-1;
 MEDLINE=96258557;PubMed=8992315 [[NCBI](#), [ExPASy](#), [EBI](#), [Israel](#), [Japan](#)]
[Lally N.C.](#), [Jenkins M.C.](#), [Dubey J.P.](#);
 "Development of a polymerase chain reaction assay for the diagnosis of neosporosis using the *Neospora caninum* 14-3-3 gene."
 Mol. Biochem. Parasitol. 75:169-178(1996).


Comments

- **SIMILARITY:** Belongs to the 14-3-3 family.

Copyright

This Swiss-Prot entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See <http://www.isb-sib.ch/announce/> or send an email to license@isb-sib.ch)

Cross-references

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
--	--------------------------	-------------------------------	----------------------------	----------------------------

Search for

NiceProt

View of

Swiss-

Prot:

P90661

[Printer-friendly view](#)
[Submit update](#)
[Quick BlastP search](#)

[\[Entry info\]](#)
[\[Name and origin\]](#)
[\[References\]](#)
[\[Comments\]](#)
[\[Cross-references\]](#)
[\[Keywords\]](#)
[\[Features\]](#)
[\[Sequence\]](#)
[\[Tools\]](#)

Note: most headings are clickable, even if they don't appear as links. They link to the [user manual](#) or [other documents](#).

Entry information

Entry name	GRA1_NEOCA
Primary accession number	P90661
Secondary accession numbers	O02507 Q25539
Entered in Swiss-Prot in	Release 36, July 1998
Sequence was last modified in	Release 36, July 1998
Annotations were last modified in	Release 44, July 2004
Name and origin of the protein	
Protein name	Dense granule protein 1 [Precursor]
Synonyms	NcDG1 Antigen Nc4.1
Gene name	Name: DG1
From	Neospora caninum [TaxID: 29176]
Taxonomy	Eukaryota ; Alveolata ; Apicomplexa ; Coccidia ; Eimeriida ; Sarcocystidae ; Neospora .

References

- [1] SEQUENCE FROM NUCLEIC ACID.
STRAIN=Nc-1;
 MEDLINE=97391130;PubMed=9247937 [[NCBI](#), [ExPASy](#), [EBI](#), [Israel](#), [Japan](#)]
[Lally N.C.](#), [Jenkins M.C.](#), [Liddell S.](#), [Dubey J.P.](#);
 "A dense granule protein (NCDG1) gene from *Neospora caninum*."
 Mol. Biochem. Parasitol. 87:239-243(1997).
- [2] SEQUENCE OF 15-217 FROM NUCLEIC ACID.
STRAIN=Nc-1;
 MEDLINE=96336081;PubMed=8705668 [[NCBI](#), [ExPASy](#), [EBI](#), [Israel](#), [Japan](#)]
[Lally N.C.](#), [Jenkins M.C.](#), [Dubey J.P.](#);
 "Evaluation of two *Neospora caninum* recombinant antigens for use in an enzyme-linked immunosorbent assay for the diagnosis of bovine neosporosis."
 Clin. Diagn. Lab. Immunol. 3:275-279(1996).

Comments

- **SUBCELLULAR LOCATION:** Secreted (*Potential*).
- **SIMILARITY:** Belongs to the Gra7 family.

Copyright

This Swiss-Prot entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See <http://www.isb-sib.ch/announce/> or send an email to license@isb-sib.ch)


Cross-references

EMBL	U72991; AAB48402.1; -. [EMBL / GenBank / DDBJ] [CoDingSequence] U82229; AAC47661.1; -. [EMBL / GenBank / DDBJ] [CoDingSequence] U36386; AAC47096.1; -. [EMBL / GenBank / DDBJ] [CoDingSequence]
InterPro	IPR008120; Gra7_protein. Graphical view of domain structure.
PRINTS	PR01747; DENSEGRNULE7.
ProDom	[Domain structure / List of seq. sharing at least 1 domain]
BLOCKS	P90661.
ProtoNet	P90661.
ProtoMap	P90661.
PRESAGE	P90661.
DIP	P90661.
ModBase	P90661.
SMR	P90661; 098B4C1B33705CC2.
SWISS-2DPAGE	Get region on 2D PAGE.
UniRef	View cluster of proteins with at least <u>50%</u> / <u>90%</u> identity.

Keywords

Antigen; Signal.

Features



[Feature table viewer](#)

Key	From	To	Length	Description
SIGNAL	<u>1</u>	<u>19</u>	19	<i>Potential.</i>
CHAIN	<u>20</u>	<u>217</u>	198	Dense granule protein 1.
CARBOHYD	<u>191</u>	<u>191</u>		N-linked (GlcNAc...) (<i>Potential</i>).
CONFLICT	<u>15</u>	<u>18</u>		GLAI -> IRHE (in Ref. <u>2</u>).

Sequence information

Length: **217 AA** [This is the length of the unprocessed precursor]

Molecular weight: **22495 Da** [This is the MW of the unprocessed precursor]

CRC64: **098B4C1B33705CC2** [This is a checksum on the sequence]

10	20	30	40	50	60
MARQATFIVA	LCVCGLAIAG	LPRLASAGDL	ATEQHEGDIG	YGVRAYAGVS	NYDGDDDAAG
70	80	90	100	110	120
NPVDSDVTD	AITDGEWPRV	VSGQKPHTTQ	KGSLIKKLAV	PVVGALTSYL	VADRVLPFLT
130	140	150	160	170	180

```
SAEEEGTESI PGKKRVKTAV GIAALVAAAA FAGLGLARTF RHFVPKKSKT VASEDSALGN
      190      200      210
SEEQYVEGTV NGSSDPEQER AGGPLIPEGD EQEVDTE
```

P90661 in [FASTA](#)
[format](#)

[View entry in original Swiss-Prot format](#)

[View entry in raw text format \(no links\)](#)

[Report form for errors/updates in this Swiss-Prot entry](#)

BLAST [BLAST submission on](#)
[ExPASy/SIB](#)
or at [NCBI \(USA\)](#)



Sequence analysis tools: [ProtParam](#), [ProtScale](#),
[Compute pI/Mw](#), [PeptideMass](#), [PeptideCutter](#),
[Dotlet \(Java\)](#)




[ScanProsite](#), [MotifScan](#)



Search the [SWISS-MODEL Repository](#)

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot				
Hosted by CBR Canada	Mirror sites:	Australia	Bolivia	China	Korea	Switzerland	Taiwan	USA

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
--	--------------------------	-------------------------------	----------------------------	----------------------------

Search for

NiceProt

View of

Swiss-

Prot:

Q25540

[Printer-friendly view](#)
[Submit update](#)
[Quick BlastP search](#)

[\[Entry info\]](#)
[\[Name and origin\]](#)
[\[References\]](#)
[\[Comments\]](#)
[\[Cross-references\]](#)
[\[Keywords\]](#)
[\[Features\]](#)
[\[Sequence\]](#)
[\[Tools\]](#)

- Note: most headings are clickable, even if they don't appear as links. They link to the [user manual](#) or [other documents](#).

Entry information

Entry name	GRA2_NEOCA
Primary accession number	Q25540
Secondary accession number	O76215
Entered in Swiss-Prot in	Release 36, July 1998
Sequence was last modified in	Release 42, October 2003
Annotations were last modified in	Release 44, July 2004
Name and origin of the protein	
Protein name	Dense granule protein 2
Synonyms	NcDG2 Antigen Nc14.1
Gene name	Name: DG2
From	Neospora caninum [TaxID: 29176]
Taxonomy	Eukaryota ; Alveolata ; Apicomplexa ; Coccidia ; Eimeriida ; Sarcocystidae ; Neospora .

References

[1] SEQUENCE FROM NUCLEIC ACID.

STRAIN=Nc-1;

MEDLINE=98324416;PubMed=9662039 [[NCBI](#), [ExPASy](#), [EBI](#), [Israel](#), [Japan](#)]

[Liddell S.](#), [Lally N.C.](#), [Jenkins M.C.](#), [Dubey J.P.](#);

"Isolation of the cDNA encoding a dense granule associated antigen (NCDG2) of *Neospora caninum*.";

Mol. Biochem. Parasitol. 93:153-158(1998).

[2] SEQUENCE OF 67-193 FROM NUCLEIC ACID.

STRAIN=Nc-1;

MEDLINE=96336081;PubMed=8705668 [[NCBI](#), [ExPASy](#), [EBI](#), [Israel](#), [Japan](#)]

[Lally N.C.](#), [Jenkins M.C.](#), [Dubey J.P.](#);

"Evaluation of two *Neospora caninum* recombinant antigens for use in an enzyme-linked immunosorbent assay for the diagnosis of bovine neosporosis.";

[Clin. Diagn. Lab. Immunol.](#) 3:275-279(1996).

Comments

- **SIMILARITY:** Belongs to the Gra6 family.

Copyright


This Swiss-Prot entry is copyright. It is produced through a collaboration between the Swiss Institute of Bioinformatics and the EMBL outstation - the European Bioinformatics Institute. There are no restrictions on its use by non-profit institutions as long as its content is in no way modified and this statement is not removed. Usage by and for commercial entities requires a license agreement (See <http://www.isb-sib.ch/announce/> or send an email to license@isb-sib.ch)

Cross-references

EMBL	AF029350; AAC39122.1; -. [EMBL / GenBank / DDBJ] [CoDingSequence] U36387; AAC47097.1; -. [EMBL / GenBank / DDBJ] [CoDingSequence]
InterPro	IPR008119 ; Gra6_protein. Graphical view of domain structure.
Pfam	PF05084 ; GRA6; 1. Pfam graphical view of domain structure.
ProDom	[Domain structure / List of seq. sharing at least 1 domain]
BLOCKS	Q25540.
ProtoNet	Q25540.
ProtoMap	Q25540.
PRESAGE	Q25540.
DIP	Q25540.
ModBase	Q25540.
SMR	Q25540 ; 7EF46F94C289BC0B.
SWISS-2DPAGE	Get region on 2D PAGE.
UniRef	View cluster of proteins with at least <u>50%</u> / <u>90%</u> identity.

Keywords

Antigen; Transmembrane.

Features

[Feature table viewer](#)



[Feature aligner](#)

Key	From	To	Length	Description
TRANSMEM	14	34	21	Potential.
TRANSMEM	153	173	21	Potential.
CARBOHYD	4	4		N-linked (GlcNAc...) (Potential).
CARBOHYD	74	74		N-linked (GlcNAc...) (Potential).
CONFLICT	67	70		GTSE -> NSAR (in Ref. 2).

Sequence information

Length: **193** Molecular weight: **20590** CRC64: **7EF46F94C289BC0B** [This is a checksum on the AA Da sequence]

10	20	30	40	50	60
MANNRTLARR	RRAFSPLTVV	MLAVTLVAFM	GVPLSSTGAA	DAADPVESVE	ANRRGYTSYG
70	80	90	100	110	120
EPPVAVGTSE	EYVNSSELAG	SRDKGNAEAE	EEAAEVETDV	QPSSVTIDTE	ERAAPSQVQV
130	140	150	160	170	180

Q Q E R M E E A D D A P K P V P V R S A V P S T V A K R Q Q A R H R V I G T A V I A A V V A A L L W K F S R R R S G A P
190
R E G G E N E N G G E E K

Q25540 in [FASTA](#)
[format](#)

[View entry in original Swiss-Prot format](#)

[View entry in raw text format \(no links\)](#)

[Report form for errors/updates in this Swiss-Prot entry](#)

BLAST [BLAST submission on](#)
[ExPASy/SIB](#)
or at [NCBI \(USA\)](#)




Sequence analysis tools: [ProtParam](#), [ProtScale](#),
[Compute pI/Mw](#), [PeptideMass](#), [PeptideCutter](#),
[Dotlet \(Java\)](#)




[ScanProsite](#), [MotifScan](#)



Search the [SWISS-MODEL Repository](#)

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot				
Hosted by CBR Canada	Mirror sites:	Australia	Bolivia	China	Korea	Switzerland	Taiwan	USA

 ExPASy Home page	Site Map	Search ExPASy	Contact us	Swiss-Prot
--	--------------------------	-------------------------------	----------------------------	----------------------------

Search for

NiceProt

View of

TrEMBL:

O61159

[Printer-friendly view](#)
[Request update](#)
[Quick BlastP search](#)

[\[Entry info\]](#)
[\[Name and origin\]](#)
[\[References\]](#)
[\[Comments\]](#)
[\[Cross-references\]](#)
[\[Keywords\]](#)
[\[Features\]](#)
[\[Sequence\]](#)
[\[Tools\]](#)

Note: most headings are clickable, even if they don't appear as links. They link to the [user manual](#) or other documents.

Entry information

Entry name	O61159
Primary accession number	O61159
Secondary accession numbers	None
Entered in TrEMBL in	Release 07, August 1998
Sequence was last modified in	Release 07, August 1998
Annotations were last modified in	Release 26, March 2004
Name and origin of the protein	
Protein name	Surface protein Nc-p43
Synonyms	None
Gene name	None
From	<u>Neospora caninum</u> [TaxID: 29176]
Taxonomy	<u>Eukaryota</u> ; <u>Alveolata</u> ; <u>Apicomplexa</u> ; <u>Coccidia</u> ; <u>Eimeriida</u> ; <u>Sarcocystidae</u> ; <u>Neospora</u> .

References

- [1] SEQUENCE FROM NUCLEIC ACID.
STRAIN=Nc-1;
 MEDLINE=96429628;PubMed=8832729 [[NCBI](#), [ExPASy](#), [EBI](#), [Israel](#), [Japan](#)]
[Hemphill A.](#), [Gottstein B.](#);
 "Identification of a major surface protein on Neospora caninum tachyzoites.";
 Parasitol. Res. 82:497-504(1996).
- [2] SEQUENCE FROM NUCLEIC ACID.
STRAIN=Nc-1;
 MEDLINE=97047711;PubMed=8926100 [[NCBI](#), [ExPASy](#), [EBI](#), [Israel](#), [Japan](#)]
[Hemphill A.](#);
 "Subcellular localization and functional characterization of Nc-p43, a major Neospora caninum tachyzoite surface protein.";
 Infect. Immun. 64:4279-4287(1996).
- [3] SEQUENCE FROM NUCLEIC ACID.
STRAIN=Nc-1;
 MEDLINE=98150127;PubMed=9488869 [[NCBI](#), [ExPASy](#), [EBI](#), [Israel](#), [Japan](#)]
[Hemphill A.](#), [Felleisen R.](#), [Connolly B.](#), [Gottstein B.](#), [Hentrich B.](#), [Muller N.](#);

"Characterization of a cDNA-clone encoding Nc-p43, a major Neospora caninum tachyzoite surface protein.";

Parasitology 115:581-590(1997).

[4] SEQUENCE FROM NUCLEIC ACID.

STRAIN=Nc-1;

Hehl A., Felleisen R., Sonda S., Fuchs N., Muller N., Hemphill A.;

Submitted (APR-1998) to the EMBL/GenBank/DDBJ databases.

Comments

None

Cross-references

EMBL U93870; AAC15250.1; -.[\[EMBL / GenBank / DDBJ\]](#) [\[CoDingSequence\]](#)

HSSP P13664; 1KZQ. [\[HSSP ENTRY / PDB\]](#)

InterPro [IPR007226](#); SAG.
[Graphical view of domain structure.](#)

Pfam [PF04092](#); SAG; 2.
[Pfam graphical view of domain structure.](#)

PRINTS [PR01801](#); SURFCEANTIGN.

ProDom [\[Domain structure / List of seq. sharing at least 1 domain\]](#)

ProtoMap [O61159](#).

PRESAGE [O61159](#).

ModBase [O61159](#).

SMR [O61159](#); CA52F48B3E7DD360.

SWISS-2DPAGE [Get region on 2D PAGE](#).

UniRef [View cluster of proteins with at least 50% / 90% identity.](#)

Keywords

None

Features

None

Sequence information

Length: **400** Molecular weight: **42051** CRC64: **CA52F48B3E7DD360** [This is a checksum on the
AA **Da** sequence]

10	20	30	40	50	60
MATHACVVR	R KADAACFAKL	SASQSCLTAK	SVNRSVSVFA	LLFGVVLAVG	VSGAPFKSEN
70	80	90	100	110	120
EKFTCLPKQG	NADQWVALVY	DSQHSITFAC	DGGRPPSKLL	SEDDGLIVCN	ESDGEDECEK
130	140	150	160	170	180
NAAPLSTFLP	GAKKEWVTGT	LQQGIKITIP	DEHYPATSKA	FRVGCKAGKN	VCLLNIVYVQS
190	200	210	220	230	240
RESEVIGQVA	HCAYSNNVRL	RPITVNPENN	GVTLICGPDG	KAFPDDYMNH	HCTELDECKE
250	260	270	280	290	300
RPYSAVFPGF	SSSFWTGEAS	GVAGATLTIP	KDQFPSTAQT	IYLGCTGHPD	DKQVTCVVPV

```
      310      320      330      340      350      360
      |       |       |       |       |       |
NIEEVAKPAG AGSNPGGGSQ PDQSSEKRDG EQVNKGKPPT GSGRGTGK LNASLNAKDK

      370      380      390      400
      |       |       |       |
GETGGENGDS PVLRGDACDE LPSYVALSAA SLTATAIFAY
```

O61159 in [FASTA](#)
[format](#)

[View entry in original TrEMBL format](#)

[View entry in raw text format \(no links\)](#)

[Request for annotation of this TrEMBL entry](#)


BLAST [BLAST submission on](#)
[ExPASy/SIB](#)
or at [NCBI \(USA\)](#)




Sequence analysis tools: [ProtParam](#), [ProtScale](#),
[Compute pI/Mw](#), [PeptideMass](#), [PeptideCutter](#),
[Dotlet \(Java\)](#)



[ScanProsite](#), [MotifScan](#)



Search the [SWISS-MODEL Repository](#)

 ExPASy Home page	Site Map	<u>Search ExPASy</u>	Contact us	Swiss-Prot				
Hosted by CBR Canada	Mirror sites:	Australia	Bolivia	China	Korea	Switzerland	Taiwan	USA

\$8.00 5 Types
 \$8.46 Estimated cost File349
 \$0.15 0.032 DialUnits File348
 \$1.70 1 Type(s) in Format 3
 \$1.70 1 Types
 \$1.85 Estimated cost File348
 \$0.02 0.006 DialUnits File10
 \$0.02 Estimated cost File10
 \$0.13 0.006 DialUnits File357
 \$6.81 3 Type(s) in Format 3
 \$6.81 3 Types
 \$6.94 Estimated cost File357
 \$0.02 0.006 DialUnits File144
 \$0.02 Estimated cost File144
 \$0.02 0.006 DialUnits File203
 \$0.02 Estimated cost File203
 \$0.02 0.006 DialUnits File358
 \$0.02 Estimated cost File358
 \$0.07 0.006 DialUnits File347
 \$0.07 Estimated cost File347
 \$0.07 0.006 DialUnits File344
 \$0.07 Estimated cost File344
 \$0.02 0.006 DialUnits File94
 \$0.02 Estimated cost File94
 \$0.02 0.006 DialUnits File65
 \$0.02 Estimated cost File65
 \$0.04 0.006 DialUnits File185
 \$0.04 Estimated cost File185
 \$0.03 0.006 DialUnits File16
 \$0.03 Estimated cost File16
 OneSearch, 16 files, 0.440 DialUnits FileOS
 \$0.24 TELNET
 \$25.70 Estimated cost this search
 \$25.70 Estimated total session cost 0.440 DialUnits

Status: Signed Off. (1 minutes)

Status: Path 1 of [Dialog Information Services via Modem]

Status: Initializing TCP/IP using (UseTelnetProto 1 ServiceID pto-dialog)
 Trying 3106000009998...Open

DIALOG INFORMATION SERVICES

PLEASE LOGON:

***** HHHHHHHH SSSSSSSS?

Status: Signing onto Dialog

ENTER PASSWORD:

***** HHHHHHHH SSSSSSSS? *****

Welcome to DIALOG

Status: Connected

Dialog level 04.11.00D

Reconnected in file OS 16aug04 16:23:39

* * * *

SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1951-2004/Aug W3

(c) format only 2004 The Dialog Corp.

*File 155: Medline has been reloaded. Accession numbers have changed. Please see HELP NEWS 154 for details.

File 5:Biosis Previews(R) 1969-2004/Aug W2

(c) 2004 BIOSIS

File 654:US Pat.Full. 1976-2004/Aug 12
 (c) Format only 2004 The Dialog Corp.
 *File 654: Reassignment data now updated monthly.
 File 349:PCT FULLTEXT 1979-2002/UB=20040812,UT=20040805
 (c) 2004 WIPO/Univentio
 File 348:EUROPEAN PATENTS 1978-2004/Aug W02
 (c) 2004 European Patent Office
 File 10:AGRICOLA 70-2004/Jun
 (c) format only 2004 The Dialog Corporation
 File 357:Derwent Biotech Res. 1982-2004/Aug W3
 (c) 2004 Thomson Derwent & ISI
 File 144:Pascal 1973-2004/Aug W2
 (c) 2004 INIST/CNRS
 File 203:AGRIS 1974-2004/Jun
 Dist by NAL, Intl Copr. All rights reserved
 File 358:Current BioTech Abs 1983-2004/Jul
 (c) 2004 DECHEMA
 File 347:JAPIO Nov 1976-2004/Apr(Updated 040802)
 (c) 2004 JPO & JAPIO
 *File 347: JAPIO data problems with year 2000 records are now fixed.
 Alerts have been run. See HELP NEWS 347 for details.
 File 344:Chinese Patents Abs Aug 1985-2004/May
 (c) 2004 European Patent Office
 File 94:JICST-EPlus 1985-2004/Jul W4
 (c)2004 Japan Science and Tech Corp(JST)
 File 65:Inside Conferences 1993-2004/Aug W2
 (c) 2004 BLDSC all rts. reserv.
 File 185:Zoological Record Online(R) 1978-2004/Jul
 (c) 2004 BIOSIS
 *File 185: File 185 has been reloaded. All accession
 numbers have changed. See HELP NEWS185.
 File 16:Gale Group PROMT(R) 1990-2004/Aug 16
 (c) 2004 The Gale Group
 *File 16: Alert feature enhanced for multiple files, duplicate
 removal, customized scheduling. See HELP ALERT.

Set Items Description

Cost is in DialUnits
 ?ds

Set	Items	Description
S1	134	NEOSPOR? (3N) (CULTUR? OR REDUC? OR CHANG? OR MODIFIC? OR - ATTEN? OR DECREAS?)
S2	107	RD (unique items)

?t s2/9/95 97 98

2/9/95 (Item 2 from file: 94)
 DIALOG(R)File 94:JICST-EPlus
 (c)2004 Japan Science and Tech Corp(JST). All rts. reserv.
 02961647 JICST ACCESSION NUMBER: 97A0890408 FILE SEGMENT: PreJICST-E
Examination on culture condition for neospora protozoa.
 ETO MARIKO (1); YAMANE ITSURO (1); KUNIYASU TAKEHIRO (1); SHIMURA KAMEO
 (1); HATAYA RYO (1); OUCHI YOSHINAO (2)
 (1) Minist. of Agric., For. and Fish., Natl. Inst. of Anim. Health; (2)
 North Dist. Livest. Hyg. Serv. Cent., Ibaraki Prefect. Gov.
 Nippon Jui Gakkai Koen Yoshishu, 1996, VOL.122nd, PAGE.119
 JOURNAL NUMBER: Z0670AAA
 LANGUAGE: Japanese COUNTRY OF PUBLICATION: Japan
 DOCUMENT TYPE: Journal
 MEDIA TYPE: Printed Publication

2/9/97 (Item 1 from file: 65)
 DIALOG(R)File 65:Inside Conferences
 (c) 2004 BLDSC all rts. reserv. All rts. reserv.

02254714 INSIDE CONFERENCE ITEM ID: CN023606104

Viability determination of culture -grown Neospora caninum

Jakubek, E.-B.; Uggla, A.; Bjoerkman, C.

CONFERENCE: Control of coccidiosis into the next millennium-International coccidiosis conference; 7th

P: 129

Newbury, Institute for Animal Health, 1997

LANGUAGE: English DOCUMENT TYPE: Conference Abstracts

CONFERENCE EDITOR(S): Shirley, M. W.; Tomley, F. M.; Freeman, B. M.

CONFERENCE SPONSOR: COST 820 Project

CONFERENCE LOCATION: Oxford

CONFERENCE DATE: Sep 1997 (199709) (199709)

BRITISH LIBRARY ITEM LOCATION: 98/07811

NOTE:

Held in conjunction with the European Union COST 820 workshop

DESCRIPTORS: coccidiosis; COST

2/9/98 (Item 2 from file: 65)

DIALOG(R)File 65:Inside Conferences

(c) 2004 BLDSC all rts. reserv. All rts. reserv.

02008929 INSIDE CONFERENCE ITEM ID: CN020883541

Evaluation of In vivo Inoculations of Culture -Derived Neospora sp. and Sarcocystis neurona Parasites for the Development of the Cyst Stage

Marsh, A. E.; Barr, B. C.; Tell, L.; Sverlow, K. W.

CONFERENCE: American Association of Veterinary Pathologists-Annual meeting; 41st

PROCEEDINGS-AMERICAN ASSOCIATION OF VETERINARY PATHOLOGISTS, 1996; CONF 41 P: 6

The Association, 1996

LANGUAGE: English DOCUMENT TYPE: Conference Abstracts and programme

CONFERENCE SPONSOR: American Association of Veterinary Pathologists

CONFERENCE LOCATION: Louisville, KY

CONFERENCE DATE: Jul 1996 (199607) (199607)

BRITISH LIBRARY ITEM LOCATION: 6622.350000

DESCRIPTORS: AAVP; veterinary pathologists

?logoff hold

16aug04 16:23:43 User228206 Session D2213.8

\$0.04 0.012 DialUnits File155

\$0.04 Estimated cost File155

\$0.07 0.012 DialUnits File5

\$0.07 Estimated cost File5

\$0.07 0.012 DialUnits File654

\$0.07 Estimated cost File654

\$0.06 0.012 DialUnits File349

\$0.06 Estimated cost File349

\$0.06 0.012 DialUnits File348

\$0.06 Estimated cost File348

\$0.03 0.012 DialUnits File10

\$0.03 Estimated cost File10

\$0.24 0.012 DialUnits File357

\$0.24 Estimated cost File357

\$0.04 0.012 DialUnits File144

\$0.04 Estimated cost File144

\$0.03 0.012 DialUnits File203

\$0.03 Estimated cost File203

\$0.05 0.012 DialUnits File358

\$0.05 Estimated cost File358

\$0.14 0.012 DialUnits File347

\$0.14 Estimated cost File347

\$0.13 0.012 DialUnits File344

\$0.13 Estimated cost File344

\$0.04 0.012 DialUnits File94

\$1.35 1 Type(s) in Format 9
 \$1.35 1 Types
\$1.39 Estimated cost File94
 \$0.14 0.037 DialUnits File65
 \$2.20 2 Type(s) in Format 9
 \$2.20 2 Types
\$2.34 Estimated cost File65
 \$0.08 0.012 DialUnits File185
\$0.08 Estimated cost File185
 \$0.07 0.012 DialUnits File16
\$0.07 Estimated cost File16
 OneSearch, 16 files, 0.224 DialUnits FileOS
\$0.24 TELNET
\$5.08 Estimated cost this search
\$5.08 Estimated total session cost 0.224 DialUnits

Status: Signed Off. (1 minutes)

SYSTEM:OS - DIALOG OneSearch

File 155:MEDLINE(R) 1951-2004/Aug W3

(c) format only 2004 The Dialog Corp.

*File 155: Medline has been reloaded. Accession numbers have changed. Please see HELP NEWS 154 for details.

File 5:Biosis Previews(R) 1969-2004/Aug W2

(c) 2004 BIOSIS

File 654:US Pat.Full. 1976-2004/Aug 12

(c) Format only 2004 The Dialog Corp.

*File 654: Reassignment data now updated monthly.

File 349:PCT FULLTEXT 1979-2002/UB=20040812,UT=20040805

(c) 2004 WIPO/Univentio

File 348:EUROPEAN PATENTS 1978-2004/Aug W02

(c) 2004 European Patent Office

File 10:AGRICOLA 70-2004/Jun

(c) format only 2004 The Dialog Corporation

File 357:Derwent Biotech Res. 1982-2004/Aug W3

(c) 2004 Thomson Derwent & ISI

File 144:Pascal 1973-2004/Aug W2

(c) 2004 INIST/CNRS

File 203:AGRIS 1974-2004/Jun

Dist by NAL, Intl Copr. All rights reserved

File 358:Current BioTech Abs 1983-2004/Jul

(c) 2004 DECHEMA

File 347:JAPIO Nov 1976-2004/Apr(Updated 040802)

(c) 2004 JPO & JAPIO

*File 347: JAPIO data problems with year 2000 records are now fixed. Alerts have been run. See HELP NEWS 347 for details.

File 344:Chinese Patents Abs Aug 1985-2004/May

(c) 2004 European Patent Office

File 94:JICST-EPlus 1985-2004/Jul W4

(c)2004 Japan Science and Tech Corp(JST)

File 65:Inside Conferences 1993-2004/Aug W2

(c) 2004 BLDSC all rts. reserv.

File 185:Zoological Record Online(R) 1978-2004/Jul

(c) 2004 BIOSIS

*File 185: File 185 has been reloaded. All accession numbers have changed. See HELP NEWS185.

File 16:Gale Group PROMT(R) 1990-2004/Aug 16

(c) 2004 The Gale Group

*File 16: Alert feature enhanced for multiple files, duplicate removal, customized scheduling. See HELP ALERT.

Set Items Description

--- -----

Cost is in DialUnits

?ds

Set Items Description

S1 134 NEOSPOR? (3N) (CULTUR? OR REDUC? OR CHANG? OR MODIFIC? OR -
ATTEN? OR DECREAS?)

S2 107 RD (unique items)

?t s2/9/9

2/9/9 (Item 9 from file: 155)

DIALOG(R)File 155:MEDLINE(R)

(c) format only 2004 The Dialog Corp. All rts. reserv.

10923841 PMID: 11068819

Humoral immune responses and safety of experimental formulations of inactivated Neospora vaccines.

Choromanski L; Block W

Bayer, Agriculture Division, Animal Health, Shawnee Mission, KS 66201, USA.

Parasitology research (GERMANY) Oct 2000, 86 (10) p851-3, ISSN 0932-0113 Journal Code: 8703571

Document type: Journal Article

Languages: ENGLISH
Main Citation Owner: NLM
Record type: Completed
Subfile: INDEX MEDICUS

Antibody titers to Neospora antigens ranged from 40 to 160 before vaccination, from 80 to 5,120 2 weeks after the first dose of vaccination, and 320 to 40,960 2 weeks after the second (booster) vaccination. A peak antibody titer of 40,960 was also detected 28 days after the booster vaccination among animals vaccinated with Neospora vaccine formulated with Bay R1005 adjuvant. In heifers inoculated with experimental formulations of Neospora vaccines, transient development of injection site reactions resulted in 1 out of 15 animals. This injection site reaction was not detectable 14 days after the first observation and measurements were made. We have also demonstrated that vaccines derived from tissue- **culture** -grown **Neospora** tachyzoites are safe and would be expected to be efficacious.

Tags: Female

Descriptors: *Antibodies, Protozoan--blood--BL; *Cattle Diseases --prevention and control--PC; *Coccidiosis--veterinary--VE; *Neospora --immunology--IM; *Protozoan Vaccines--immunology--IM; Animals; Cattle; Vaccination

CAS Registry No.: 0 (Antibodies, Protozoan); 0 (Protozoan Vaccines)

Record Date Created: 20010227

Record Date Completed: 20010301

?t s2/3,kwic/19 21 28 29 30 36 48 49 56 58 59 70 71 74 75

2/3,KWIC/19 (Item 1 from file: 5)

DIALOG(R)File 5:Biosis Previews(R)

(c) 2004 BIOSIS. All rts. reserv.

0014868771 BIOSIS NO.: 200400249528

Recombinant neospora antigens and their uses

AUTHOR: Conrad Patricia A (Reprint); Barr Bradd C; Anderson Mark L; Sverlow Karen W

AUTHOR ADDRESS: Davis, CA, USA**USA

JOURNAL: Official Gazette of the United States Patent and Trademark Office
Patents 1281 (1): Apr. 6, 2004 2004

MEDIUM: e-file

PATENT NUMBER: US 6716423 PATENT DATE GRANTED: April 06, 2004 20040406

PATENT CLASSIFICATION: 424-931 PATENT ASSIGNEE: The Regents of the
University of California PATENT COUNTRY: USA

ISSN: 0098-1133 (ISSN print)

DOCUMENT TYPE: Patent

RECORD TYPE: Abstract

LANGUAGE: English

ABSTRACT: The present invention provides isolated bovine **Neospora cultures** . Also provided are recombinant immunodominant **Neospora** antigens. The **cultures** and antigens are used to develop diagnostic assays for the detection of Neospora infections in...

2/3,KWIC/21 (Item 3 from file: 5)

DIALOG(R)File 5:Biosis Previews(R)

(c) 2004 BIOSIS. All rts. reserv.

0014412576 BIOSIS NO.: 200300371295

Adaptation in cell- culture of Neospora caninum isolates to changing environments: An argument in favor of its asexual propagation.

AUTHOR: Perez-Zaballos F J (Reprint); Costas E (Reprint); Alvarez-Garcia G (Reprint); Collantes-Fernandez E (Reprint); Ortega-Mora L M (Reprint)

AUTHOR ADDRESS: Departamento Sanidad Animal, Facultad de Veterinaria,
Universidad Complutense de Madrid, Ciudad Universitaria s/n, 28040,
Madrid, Spain**Spain

JOURNAL: Infection Genetics and Evolution 2 (4): p303 April 2003 2003

MEDIUM: print

CONFERENCE/MEETING: Sixth International Meeting on Molecular Epidemiology
and Evolutionary Genetics of Infectious Diseases Paris, France July 23,

2002-July 27, 200320020723
ISSN: 1567-1348 (ISSN print)
DOCUMENT TYPE: Meeting; Meeting Abstract
RECORD TYPE: Citation
LANGUAGE: English

Adaptation in cell- culture of Neospora caninum isolates to changing environments: An argument in favor of its asexual propagation.

2/3,KWIC/28 (Item 2 from file: 654)
DIALOG(R)File 654:US Pat.Full.
(c) Format only 2004 The Dialog Corp. All rts. reserv.

5597804 **IMAGE Available
Derwent Accession: 2000-664889

Utility

Recombinant neospora antigens and their uses

Inventor: Conrad, Patricia A., Davis, CA
Barr, Bradd C., Davis, CA
Anderson, Mark L., Davis, CA
Sverlow, Karen W., Vacaville, CA
Assignee: The Regents of the University of California (02), Oakland, CA
Examiner: Navarro, Mark (Art Unit: 165)
Law Firm: Townsend and Townsend and Crew L.L.P.

	Publication Number	Kind	Date	Application Number	Filing Date
Main Patent	US 6716423	A	20040406	US 2000612858	20000710
Continuation	US 6376196	A		US 99281766	19990330
CIP	US 5889166	A	19990330	US 96645951	19960510
CIP	US 5707617	A	19980113	US 94327516	19941020
CIP	Abandoned			US 94215858	19940321

Fulltext Word Count: 19836

Abstract:

The present invention provides isolated bovine **Neospora cultures**. Also provided are recombinant immunodominant **Neospora** antigens. The **cultures** and antigens are used to develop diagnostic assays for the detection of **Neospora** infections in...

Description of the Invention:

...A "biologically pure bovine **Neospora culture**" refers to a continuous in vitro **culture** of bovine **Neospora** organisms (e.g. tachyzoites) which is substantially free of other organisms other than the host... **Neospora** tachyzoite **cultures** of the invention have been deposited with the American Type Culture Collection, Rockville, Md. on... antibodies raised against the tachyzoite of interest. By using these methods, immunodominant antigens from different **cultures** or species of **Neospora** can be identified...**Neospora** antigens. If isolated proteins are used, they may be recombinantly produced or isolated from **Neospora cultures**. Synthetic peptides made using the protein sequences may also be used...combinations of capture agent and labeled binding agent can be used. For instance, an isolated **Neospora** protein or **culture** can be used as the capture agent and labeled anti-bovine antibodies specific for the...The present invention provides biologically pure **cultures** of bovine **Neospora**. Two such **cultures** have been deposited with the ATCC and given ATCC Accession No. 75710 (BPA1), and ATCC...An **attenuated Neospora** vaccine can only be used in the absence of a risk of human infection should...Collection #CCL209) was found to be the best cell monolayer for the cultivation of bovine **Neospora**. One of these **cultures** (BPA6) has been shown to induce bradyzoite cysts in mice... these cows could be tested repeatedly over a 6 to 12 month period to determine **changes** in the **Neospora** titer. Peak titers of 640 to 2,560 were apparent within the first 20 daysThis example describes experimental

infections of pregnant cows with **culture** -derived **Neospora** tachyzoites ...In addition, polyclonal monospecific antisera to clones N54 and N57 only bound to **Neospora** antigens on **reducing** and nonreducing western blots. Rabbit anti-N54 recognized Neospora bands of molecular weights 97.2...

Non-exemplary or Dependent Claim(s):

- ...4. The method of claim 1, wherein the isolated bovine **Neospora** tachyzoite is **attenuated** .
- ...9. The method of claim 6, wherein the isolated bovine **Neospora** tachyzoite is **attenuated** .
- ...composition comprising a pharmaceutically acceptable carrier and an immunogenically effective amount of an isolated bovine **Neospora** tachyzoite that is **attenuated** , whereby the bovine animal is protected from abortion induced by Neospora infection...

2/3,KWIC/29 (Item 3 from file: 654)

DIALOG(R) File 654:US Pat.Full.

(c) Format only 2004 The Dialog Corp. All rts. reserv.

0005594082

Derwent Accession: 1998-258985

Attenuated **live** neospora **vaccine**

Inventor: Brake, David, INV

Blagburn, Byron, INV

Lindsay, David, INV

Assignee: PFIZER INC (02), NEW YORK, NY

Correspondence Address: Dr. Peter C. Richardson Pfizer Inc., 235 East 42nd Street, New York, NY, 10017-5755, US

	Publication Number	Kind	Date	Application Number	Filing Date
Main Patent	US 20040062772	A1	20040401	US 2003669941	20030924
Continuation	US 6656479			US 2001952388	20010912
Continuation	ABANDONED			US 99260414	19990226
Continuation	ABANDONED			US 97967744	19971110
Provisional				US 60-31248	19961112

Fulltext Word Count: 11700

Attenuated **live** neospora **vaccine**

Summary of the Invention:

...invention relates to attenuated strains of the pathogenic protozoan, Neospora, and to live vaccines against **neosporosis** prepared from the **attenuated** strains which are useful in the prevention of clinical disease and abortion in mammals...0005] WO 9525541 discloses a biologically pure **culture** of bovine **Neospora** , methods of detecting anti-Neospora antibodies and Neospora-specific nucleic acids, and a composition containing...

...Neospora antigen and carrier for use as a vaccine. WO 9525541 does not, however, teach **attenuated** live **cultures** of **Neospora** , or live vaccines prepared therefrom which are able to trigger a protective immune response in...of cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated**

pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an

Description of the Invention:

...0109] The results indicate that a formulation comprising **attenuated** , live **Neospora** tachyzoites and an adjuvant is at least as effective and safe for use as a...0114] Table 20 demonstrates the ability of a vaccine comprising live, **attenuated** tachyzoites of **Neospora** to protect pygmy goat does against Neospora-induced abortion. All 4 goat does vaccinated with...

Exemplary or Independent Claim(s):

...of cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

2/3,KWIC/30 (Item 4 from file: 654)

DIALOG(R)File 654:US Pat.Full.

(c) Format only 2004 The Dialog Corp. All rts. reserv.

5451651

Derwent Accession: 1998-258985

Utility

C/ Attenuated **live** neospora vaccine
; PROTECTS A MAMMAL **AGAINST NEOSPOROSIS**

Inventor: Brake, David A, East Lyme, CT
Blagburn, Byron L, Auburn, AL
Lindsay, David S, Christiansburg, VA

Assignee: Pfizer Inc. (02), New York, NY
Pfizer Inc (Code: 65376)

Examiner: Smith, Lynette R. F. (Art Unit: 165)

Assistant Examiner: Portner, Ginny Allen

Law Firm: Scully, Scott, Murphy & Presser

	Publication Number	Kind	Date	Application Number	Filing Date
Main Patent	US 6656479	A	20031202	US 2001952388	20010912
Continuation	Abandoned			US 99260414	19990226
Continuation	Abandoned			US 97967744	19971110

Fulltext Word Count: 9627

Attenuated **live** neospora vaccine...

Summary of the Invention:

...invention relates to attenuated strains of the pathogenic protozoan, Neospora, and to live vaccines against **neosporosis** prepared from the **attenuated** strains which are useful in the prevention of clinical

disease and abortion in mammals...WO 9525541 discloses a biologically pure **culture** of bovine **Neospora**, methods of detecting anti-**Neospora** antibodies and **Neospora**-specific nucleic acids, and a composition containing...

...**Neospora** antigen and carrier for use as a vaccine. WO 9525541 does not, however, teach **attenuated** live **cultures** of **Neospora**, or live vaccines prepared therefrom which are able to trigger a protective immune response in...

...of cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...Applicants have discovered that cells of a pathogenic strain of a species of **Neospora** may be **attenuated**, and that the resulting attenuated cells are capable of ...of cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...The term "attenuated" as used herein describes a cell, **culture**, or strain of **Neospora** exhibiting a detectable **reduction** in infectivity or virulence in vitro or in vivo as compared to that of the parent strain of **Neospora** from which the **attenuated** cell, culture, or strain is derived. Reduction in virulence encompasses any detectable decrease in any...Preparation of **Attenuated** Strains of **Neospora** ...Since the invention is based on the discovery that cells of a pathogenic strain of **Neospora** may be **attenuated**, and that the resulting attenuated cells are capable of triggering an immune response that protects...exposure, is preferably that amount which results in producing one or more viable cells of **Neospora** that exhibit an **attenuated** level of pathogenicity but that are capable of triggering an immune response that protects against...

...Pathogenic strains of **Neospora** may also be **attenuated** using recombinant DNA technology according to techniques known in the art, and the present inventionAn **attenuated** strain of **Neospora** may be derived from any pathogenic strain of any species of the genus including, but...

...Both parental strains and **attenuated** strains of **Neospora** may be **cultured** in vitro by infecting any receptive cell line, preferably a mammalian cell line, with tachyzoites...

...according to known techniques described in the art. Mammalian cell lines in which tachyzoites of **Neospora** can be **cultured** include, ... Mammalian cell cultures can be grown, and cell **cultures** infected with **Neospora** can be maintained, in any one of several culture media described in the art. For...

...but in which the fetal bovine serum is increased to 10% (v/v) (growth medium). **Attenuated** strains of **Neospora** having novel auxotrophies will require appropriate modification to the culture medium to support growth, as...

... **Neospora** -infected monolayer **cultures** of mammalian cells are typically maintained under standard tissue culture conditions such as, for example...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...The vaccine of the invention comprises live cells of an **attenuated** strain of **Neospora** , either as the sole antigenic component or in combination with one or more other antigens...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...formulations. Immunomodulatory agents are selected based on their ability to maintain both viability of the **attenuated Neospora** cells and ability of the cells to trigger a protective immune response in the vaccinated...term "protection" is used broadly and is not limited to absolute prevention of infection by **Neospora** , but includes a **reduction** in infectivity, or in the severity of a disease or condition resulting from infection, including...

Description of the Invention:

...The results indicate that a formulation comprising **attenuated** , live **Neospora** tachyzoites and an adjuvant is at ...Table 20 demonstrates the ability of a vaccine comprising live, **attenuated** tachyzoites of **Neospora** to protect pygmy goat does against Neospora-induced abortion. All 4 goat does vaccinated with...

2/3,KWIC/36 (Item 10 from file: 654)

DIALOG(R) File 654:US Pat.Full.

(c) Format only 2004 The Dialog Corp. All rts. reserv.

0005091678

Derwent Accession: 1999-288171

Neospora vaccines

Inventor: Leszek Choromanski, INV

Karen Brown, INV

Correspondence Address: INTERVET INC, 405 STATE STREET PO BOX 318, MILLSBORO, DE, 19966, US

	Publication Number	Kind	Date	Application Number	Filing Date
Main Patent	US 20020146436	A1	20021010	US 2002115478	20020402
Division	PENDING			US 97954531	19971020

Fulltext Word Count: 6469

Abstract:

A Neospora caninum vaccine comprising tissue **culture** grown **Neospora** and methods of making and using said vaccines. Neospora caninum vaccines described include those containing...

Summary of the Invention:

...this invention to describe a vaccine composition for protection of mammals from disease caused by **Neospora** caninum comprising tissue **culture** grown **Neospora** caninum tachyzoites as a whole culture or in an extract form or as subunit antigens...

...in a susceptible tissue culture until a cytopathic effect (CPE) is produced, harvesting said tissue **culture** grown **Neospora** caninum and formulating said harvest into a vaccine. A modified live vaccine produced in this...

...caninum in a susceptible tissue culture until a CPE is produced; 2) harvesting said tissue **culture** grown **Neospora** caninum; 3) inactivating said harvested tissue **culture** grown **Neospora** caninum; and 4) adjuvanting the inactivated harvested tissue **culture** grown **Neospora** caninum to produce a vaccine. Still another method of producing a vaccine for protection of...

...caninum in a susceptible tissue culture until a CPE is produced; 2) harvesting said tissue **culture** grown **Neospora** caninum; 3) extracting protective antigens from the harvested tissue **culture** grown **Neospora** caninum to produce subunits; 4) inactivating the subunits if necessary; and 5) adjuvanting the subunits...sup]6 TCID[sub]50/mL are produced. A Master Seed means that the tissue **culture** grown **Neospora** sp is grown to a high titer, aliquoted into equivalent volumes in freezing vials and ...mutated **Neospora** caninum in a susceptible tissue culture until CPE is produced, harvesting the tissue **culture** grown **Neospora** sp and formulating said harvest into a vaccine. Formulation may include addition of stabilizers and...

...**Neospora** caninum in a susceptible tissue culture until a CPE is produced, harvesting said tissue **culture** grown **Neospora** caninum, inactivating said harvested tissue **culture** grown **Neospora** caninum; and adjuvanting the inactivated harvested tissue **culture** grown **Neospora** caninum to produce a vaccine a susceptible tissue culture until a CPE is produced, harvesting said tissue **culture** grown **Neospora** caninum, extracting protective antigens from the harvested tissue **culture** grown **Neospora** caninum to produce protective antigen subunits, inactivating the subunits if necessary; and adjuvanting the subunits...

Description of the Invention:

...of DMEM. Each roller bottle contained 264 mL of DMEM plus 10% Horse Serum. The **neospora** -infected tissue **cultures** were incubated at 37 degree C. until at least 50% of the cells demonstrated CPE...

Exemplary or Independent Claim(s):

1. A **Neospora** caninum vaccine comprising tissue **culture** grown **Neospora** .

...

...in a susceptible tissue culture until a cytopathic effect is produced; b. harvesting said tissue **culture** grown **Neospora** caninum; and c. formulating said harvest into a vaccine...

...in a susceptible tissue culture until a cytopathic effect is produced; b. harvesting said tissue **culture** grown **Neospora** caninum; c. inactivating said harvested tissue **culture** grown **Neospora** caninum; and d. adjuvanting the inactivated harvested tissue **culture** grown **Neospora** caninum to produce a vaccine...

...in a susceptible tissue culture until a cytopathic effect is produced; b. harvesting said tissue **culture** grown **Neospora** caninum; c. extracting one or more protective antigens from the harvested tissue **culture** grown **Neospora** caninum to produce a subunit; d. inactivating the subunit(s), optionally; and e. adjuvanting the...

...in a susceptible tissue culture until a cytopathic effect is produced; b. harvesting said tissue **culture** grown **Neospora** caninum; c. inactivating the **Neospora** caninum harvest; d. extracting one or more protective antigens from the harvested, inactivated tissue **culture** grown **Neospora** caninum to produce a subunit(s); and e. adjuvanting the subunit(s) to produce a...

Non-exemplary or Dependent Claim(s):

2. The **Neospora** caninum vaccine according to claim 1 wherein the tissue **culture** grown **Neospora** comprises an antigen selected from the group consisting of a whole **culture** of **Neospora** tachyzoites, an inactivated tissue **culture** of **Neospora** tachyzoites, a modified

live tissue **culture** **Neospora** tachyzoites, an extract from
Neospora tachyzoites and one or more subunits obtained from Neospora
tachyzoites...

2/3,KWIC/48 (Item 1 from file: 349)
DIALOG(R)File 349:PCT FULLTEXT
(c) 2004 WIPO/Univentio. All rts. reserv.

01106055

LIVE ANTENUATED PARASITE VACCINE
VACCIN DE PARASITE VIVANT ATTENUE

Patent Applicant/Assignee:

AKZO NOBEL N V, Velperweg 76, NL-6824 BM Arnhem, NL, NL (Residence), NL
(Nationality), (For all designated states except: US)

Patent Applicant/Inventor:

VAN POPPEL Nicole Francisca Johanna, Van Welderenstraat 105 A, NL-6511 MG
Nijmegen, NL, NL (Residence), NL (Nationality), (Designated only for:
US)

VERMEULEN Arnoldus Nicolaas, Korhoenderveld 34, NL-5431 HH Cuyk, NL, NL
(Residence), NL (Nationality), (Designated only for: US)

SCHAAP Theodorus Cornelis, Van de Does de Willeboissingel 53, NL-5211
CE's-Hertogenbosch, NL, NL (Residence), NL (Nationality), (Designated
only for: US)

Legal Representative:

MESTROM J J L (et al) (agent), Intervet International B.V., P.O. Box 31,
NL-5830 AA Boxmeer, NL,

Patent and Priority Information (Country, Number, Date):

Patent: WO 200426903 A2-A3 20040401 (WO 0426903)

Application: WO 2003EP10696 20030919 (PCT/WO EP03010696)

Priority Application: EP 200278953 20020920

Designated States:

(Protection type is "patent" unless otherwise stated - for applications
prior to 2004)

AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK DM DZ

EC EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR

LS LT LU LV MA MD MG MK MN MW MX MZ NI NO NZ OM PG PH PL PT RO RU SC SD

SE SG SK SL SY TJ TM TN TR TT TZ UA UG US UZ VC VN YU ZA ZM ZW

(EP) AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL PT RO SE
SI SK TR

(OA) BF BJ CF CG CI CM GA GN GQ GW ML MR NE SN TD TG

(AP) GH GM KE LS MW MZ SD SL SZ TZ UG ZM ZW

(EA) AM AZ BY KG KZ MD RU TJ TM

Publication Language: English

Filing Language: English

Fulltext Word Count: 12092

Fulltext Availability:

Claims

Claim

... 3, characterised in that said parasite belongs to the genus Eimeria,
Cryptosporidium, Toxoplasma, Sarcocystis or **Neospora** . 5) **Attenuated**
live parasite according to claim 2, characterised in that said parasite
belongs to the family...

2/3,KWIC/49 (Item 2 from file: 349)
DIALOG(R)File 349:PCT FULLTEXT
(c) 2004 WIPO/Univentio. All rts. reserv.

01008473 **Image available**

SEROLOGICAL ASSAY FOR NEOSPOA CANINUM
DOSAGE SEROLOGIQUE DE NEOSPOA CANINUM

Patent Applicant/Assignee:

PFIZER PRODUCTS INC, Eastern Point Road, Groton, CT 06340, US, US

(Residence), US (Nationality), (For all designated states except: US)

Patent Applicant/Inventor:

BRAKE David Alan, Pfizer Global Research and Development, Eastern Point
Road, Groton, CT 06340, US, US (Residence), US (Nationality),
(Designated only for: US)

RITTER Dianne Marie, Pfizer Global Research and Development, Eastern
Point Road, Groton, CT 06340, US, US (Residence), US (Nationality),
(Designated only for: US)

Legal Representative:

LUMB J Trevor (et al) (agent), c/o Simpson, Alison Urquhart-Dykes & Lord,
30 Welbeck Street, London W1G 8ER, GB,

Patent and Priority Information (Country, Number, Date):

Patent: WO 200338434 A2-A3 20030508 (WO 0338434)

Application: WO 2002IB4125 20021008 (PCT/WO IB0204125)

Priority Application: US 2001334811 20011031

Designated States:

(Protection type is "patent" unless otherwise stated - for applications
prior to 2004)

AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK DM DZ
EC EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR
LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ OM PH PL PT RO RU SD SE SG SI
SK SL TJ TM TN TR TT TZ UA UG US UZ VN YU ZA ZM ZW

(EP) AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LU MC NL PT SE SK TR

(OA) BF BJ CF CG CI CM GA GN GQ GW ML MR NE SN TD TG

(AP) GH GM KE LS MW MZ SD SL SZ TZ UG ZM ZW

(EA) AM AZ BY KG KZ MD RU TJ TM

Publication Language: English

Filing Language: English

Fulltext Word Count: 2938

Fulltext Availability:

Detailed Description

Detailed Description

... with Neospora-based vaccines (see, e.g. U.S. Serial 5 No. 09/260,414 "
Attenuated Live Neospora Vaccine", incorporated herein by reference
and ...vaccinated on day 0 and day 21 with either a tachyzoite-based i)
modified-live, **attenuated Neospora** vaccine (USPTO patent application
serial 091260,414 "**Attenuated Live Neospora Vaccine**", incorporated
herein by reference) or ii) inactivated, whole cell homogenate Neospora
vaccine (USPTO patent...

2/3,KWIC/56 (Item 9 from file: 349)

DIALOG(R)File 349:PCT FULLTEXT

(c) 2004 WIPO/Univentio. All rts. reserv.

00488951

NEOSPORA VACCINES

VACCINS NEOSPORA

Patent Applicant/Assignee:

BAYER CORPORATION,

Inventor(s):

CHOROMANSKI Leszek,

BROWN Karen K,

Patent and Priority Information (Country, Number, Date):

Patent: WO 9920303 A1 19990429

Application: WO 98US21515 19981013 (PCT/WO US9821515)

Priority Application: US 97954531 19971020

Designated States:

(Protection type is "patent" unless otherwise stated - for applications
prior to 2004)

AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GD GE GH
GM HR HU ID IL IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD MG MK MN MW
MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT UA UG UZ VN YU ZW GH
GM KE LS MW SD SZ UG ZW AM AZ BY KG KZ MD RU TJ TM AT BE CH CY DE DK ES
FI FR GB GR IE IT LU MC NL PT SE BF BJ CF CG CI CM GA GN GW ML MR NE SN
TD TG

Publication Language: English
Fulltext Word Count: 6025

Fulltext Availability:
Detailed Description
Claims

English Abstract

A i(Neospora caninum) vaccine comprising tissue **culture** grown **Neospora** and methods of making and using said vaccines. i(Neospora caninum) vaccines described include those...

French Abstract

L'invention concerne un vaccin appele i(Neospora caninum) comprenant **Neospora** obtenue par **culture** tissulaire, et des procedes de production et d'utilisation desdits vaccins. Les vaccins i(Neospora...

Detailed Description

- ... this invention to describe a vaccine composition for protection of mammals from disease caused by **Neospora** caninum comprising tissue **culture** grown **Neospora** caninum tachyzoites as a whole culture or in an extract form or as subunit antigens...
- ...in a susceptible tissue culture until a cytopathic effect (CPE) is produced, harvesting said tissue **culture** grown **Neospora** caninum and formulating said harvest into a vaccine. A modified live vaccine produced in this...
- ...caninum in a susceptible tissue culture until a CPE is produced; 2) harvesting said tissue **culture** grown **Neospora** caninum; 3) inactivating said harvested tissue **culture** grown **Neospora** caninum; and 4) adjuvanting the inactivated harvested tissue **culture** grown **Neospora** caninum to produce a vaccine. Still another method of producing a vaccine for protection of...
- ...caninum in a susceptible tissue culture until a CPE is produced; 2) harvesting said tissue **culture** grown **Neospora** caninum; 3) extracting protective antigens from the harvested tissue **culture** grown **Neospora** caninum to produce subunits; 4) inactivating the subunits if necessary; and 5) adjuvanting the subunits...and more preferably, jX106 TCID50/mL are produced. A Master Seed means that the tissue **culture** grown **Neospora** sp is grown to a high titer, aliquoted into equivalent volumes in freezing vials and...
- ...mutated **Neospora** caninum in a susceptible tissue culture until CPE is produced, harvesting the tissue **culture** grown **Neospora** sp and formulating said harvest into a vaccine.

Formulation may include addition of stabilizers and...

- ...**Neospora** caninum in a susceptible tissue culture until a CPE is produced, harvesting said tissue **culture** grown **Neospora** caninum, inactivating said harvested tissue **culture** grown **Neospora** caninum; and adjuvanting the inactivated harvested tissue **culture** grown **Neospora** caninum ...**Neospora** caninum in a susceptible tissue culture until a CPE is produced, harvesting said tissue **culture** grown **Neospora** caninum, extracting protective antigens from the harvested tissue **culture** grown **Neospora** caninum to produce protective antigen subunits, inactivating the subunits if necessary; and adjuvanting the subunits...of DMEM. Each roller bottle contained 264 mL of DMEM plus 10% Horse Serum. The **neospora** -infected tissue **cultures** were

incubated at 37°C until at least 50% of the cells demonstrated CPE
(approximately 7...

Claim

1. A *Neospora caninum* vaccine comprising tissue culture grown **Neospora**.

2 The *Neospora caninum* vaccine according to Claim 1 wherein the tissue culture grown **Neospora** comprises an antigen selected from the group consisting of a whole culture of **Neospora** tachyzoites, an inactivated tissue culture of **Neospora** tachyzoites, a modified live tissue culture **Neospora** tachyzoites, an extract from *Neospora* tachyzoites and one or more subunits obtained from *Neospora* tachyzoites...

...in a susceptible tissue culture until a cytopathic effect is produced;
b. harvesting said tissue culture grown **Neospora caninum**; and
c. formulating said harvest into a vaccine.

11 A method of producing a...

...in a susceptible tissue culture until a cytopathic effect is produced;
b. harvesting said tissue culture grown **Neospora caninum**;
c. inactivating said harvested tissue culture grown **Neospora caninum**; and
d. adjuvanting the inactivated harvested tissue culture grown **Neospora caninum** to produce a vaccine.

12 A method of producing a *Neospora* subunit vaccine comprising...

...in a susceptible tissue culture until a cytopathic effect is produced;
b. harvesting said tissue culture grown **Neospora caninum**;
c. extracting one or more protective antigens from the harvested tissue culture grown **Neospora caninum** to produce a subunit;
d. inactivating the subunit(s), optionally; and
- 28
e...

...in a susceptible tissue culture until a cytopathic effect is produced;
b. harvesting said tissue culture grown **Neospora caninum**;
c. inactivating the *Neospora caninum* harvest;
d. extracting one or more protective antigens from the harvested, inactivated tissue culture grown **Neospora caninum** to produce

2/3, KWIC/58 (Item 11 from file: 349)
DIALOG(R) File 349: PCT FULLTEXT
(c) 2004 WIPO/Univentio. All rights reserved.

00376339

A VACCINE AGAINST TOXOPLASMA GONDII
VACCIN CONTRE TOXOPLASMA GONDII

Patent Applicant/Assignee:

TRUSTEES OF DARTMOUTH COLLEGE,
KASPER Lloyd H,
KHAN Imtiaz A,
MATSUURA Tadashi,

Inventor(s):

KASPER Lloyd H,

KHAN Imtiaz A,
MATSUURA Tadashi,
Patent and Priority Information (Country, Number, Date):
Patent: WO 9717082 A1 19970515
Application: WO 96US18100 19961108 (PCT/WO US9618100)
Priority Application: US 956513 19951109
Designated States:
(Protection type is "patent" unless otherwise stated - for applications
prior to 2004)
AU CA JP US AT BE CH DE DK ES FI FR GB GR IE IT LU MC NL PT SE
Publication Language: English
Fulltext Word Count: 2188

Fulltext Availability:
Claims

Claim

... Neospora antigen.

2 The vaccine of claim 1 wherein the Neospora antigen
5 comprises live, **attenuated Neospora** .

3 The vaccine of claim 2 further comprising a
pharmaceutically acceptable carrier.

4 The vaccine...

...The method of claim 6 wherein the Neospora antigen
of the administered vaccine comprises live, **attenuated Neospora**

8 The method of claim 7 wherein the vaccine further
comprises a pharmaceutically acceptable carrier...

2/3,KWIC/59 (Item 12 from file: 349)
DIALOG(R)File 349:PCT FULLTEXT
(c) 2004 WIPO/Univentio. All rts. reserv.

00307389

BOVINE NEOSPORA ISOLATES AND THEIR USES
ISOLATS DE NEOSPORA BOVIN ET LEUR UTILISATION

Patent Applicant/Assignee:

THE REGENTS OF THE UNIVERSITY OF CALIFORNIA,
Inventor(s):

CONRAD Patricia A,
BARR Bradd C,
ANDERSON Mark L,
SVERLOW Karen W,

Patent and Priority Information (Country, Number, Date):

Patent: WO 9525541 A1 19950928
Application: WO 95US3174 19950314 (PCT/WO US9503174)
Priority Application: US 94215858 19940321; US 94327516 19941020

Designated States:

(Protection type is "patent" unless otherwise stated - for applications
prior to 2004)

AU CA JP NZ AT BE CH DE DK ES FR GB GR IE IT LU MC NL PT SE

Publication Language: English

Fulltext Word Count: 15763

Fulltext Availability:

Detailed Description
Claims

English Abstract

The present invention provides isolated bovine **Neospora cultures** . The
cultures are used to develop diagnostic assays for the detection of
Neospora infections in cattle and...

French Abstract

La presente invention concerne des **cultures** isolees de **Neospora** bovin. Ces **cultures** sont utilisees pour mettre au point des dosages diagnostiques en vue de la detection d...

Detailed Description

... blood, serum, plasma, urine,
ascites fluid, cerebrospinal fluid and fetal fluid.

A "biologically pure bovine **Neospora culture** " refers to a continuous in vitro culture of bovine **Neospora** organisms (e.g. tachyzoites...from the cultures of the invention.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides **Neospora cultures** isolated from cattle. The cultures are useful in a variety of applications including the production...

...proteins for diagnostic assays and the preparation of immunogenic proteins for use in vaccine compositions.

Neospora tachyzoite **cultures** of the invention have been deposited with the American Type Culture Collection, Rockville, Maryland on...antigens. if 3S isolated proteins are used, they may be recombinantly produced or isolated from **Neospora cultures**. Synthetic peptides made using the protein sequences may also be used.

Methods of production of...combinations of capture agent and labelled binding agent can be used. For instance, an isolated **Neospora** protein or **culture** can be used as the capture agent and labelled anti-bovine antibodies specific for the...to either cattle or the definitive host can comprise tachyzoite and/or bradyzoite antigens.

An **attenuated Neospora** vaccine can only be used in the absence of a risk of human infection...CCL209) was found to be the best cell monolayer for the cultivation of bovine 35 **Neospora**. One of these **cultures** (BPA6) has been shown to induce bradyzoite cysts in mice.

MATERIALS AND METHODS

Pathological examination...these cows could be tested repeatedly over a 6 to 12 month period to determine **changes** in the **Neospora** titer.

Peak titers of 640 to 2,560 were apparent within the first 20 days...the Toxoplasma amplification product.

Example 5

This example describes experimental infections of pregnant cows with **culture**-derived **Neospora** tachyzoites.

Three cows were inoculated with 8×10^6 tachyzoites of the BPA1 **Neospora** isolate...

Claim

1 A biologically pure **culture** of bovine **Neospora**.

2 The **culture** of claim 1, which is selected
from the group consisting of ATCC Accession No. (BPA1...

...7 The method of claim 3, wherein the antigen
is a protein isolated from a **Neospora culture** .

8 The method of claim 7, the isolated
protein has a molecular weight of 106...

2/3,KWIC/70 (Item 10 from file: 348)
DIALOG(R) File 348:EUROPEAN PATENTS
(c) 2004 European Patent Office. All rts. reserv.

00923338

Attenuated **live neospora vaccine**
Attenuierter **lebender Neospora Impfstoff**
Neospora vaccin vivant atténue

PATENT ASSIGNEE:

PFIZER INC., (200962), 235 East 42nd Street, New York, N.Y. 10017-5755,
(US), (Proprietor designated states: all)
AUBURN UNIVERSITY, (1311211), 309 Samford Hall, Auburn, AL 36849-5112,
(US), (Proprietor designated states: all)

INVENTOR:

Brake, David, 196 Upper Pattagansett Road, East Lyme, Connecticut 0633,
(US)
Blagburn, Byron, 309 Watercrest Court, Auburn, Alabama 36830, (US)
Lindsay, David, 1655 Sleepy Hollow Road, Christiansburg, Virginia 24073,
(US)

LEGAL REPRESENTATIVE:

Green, Mark Charles (57752), Urquhart-Dykes & Lord LLP 30 Welbeck Street,
London W1G 8ER, (GB)

PATENT (CC, No, Kind, Date): EP 841392 A2 980513 (Basic)
EP 841392 A3 990324
EP 841392 B1 040714

APPLICATION (CC, No, Date): EP 97308133 971014;

PRIORITY (CC, No, Date): US 31248 P 961112

DESIGNATED STATES: AT; BE; CH; DE; DK; ES; FI; FR; GB; GR; IE; IT; LI; LU;
NL; PT; SE

INTERNATIONAL PATENT CLASS: C12N-001/10; A61K-039/002; A61K-039/00;

C12N-001/10; C12R-1:90

ABSTRACT WORD COUNT: 34

LANGUAGE (Publication,Procedural,Application): English; English; English

FULLTEXT AVAILABILITY:

Available Text	Language	Update	Word Count
CLAIMS A	(English)	199820	527
CLAIMS B	(English)	200429	529
CLAIMS B	(German)	200429	522
CLAIMS B	(French)	200429	634
SPEC A	(English)	199820	8643
SPEC B	(English)	200429	8493
Total word count - document A			9173
Total word count - document B			10178
Total word count - documents A + B			19351

Attenuated **live neospora vaccine**
Attenuierter **lebender Neospora Impfstoff**
Neospora vaccin vivant atténue

...SPECIFICATION invention relates to attenuated strains of the pathogenic
protozoan, Neospora, and to live vaccines against **neosporosis** prepared
from the **attenuated** strains which are useful in the prevention of
clinical disease and abortion in mammals.

BACKGROUND...

...al., 1995, Am. J. Vet. Res. 56:1176-1180.

WO 9525541 discloses a biologically pure **culture** of bovine **Neospora**, methods of detecting anti-Neospora antibodies and Neospora-specific nucleic acids, and a composition containing...

...Neospora antigen and carrier for use as a vaccine. WO 9525541 does not, however, teach **attenuated** live **cultures** of **Neospora**, or live vaccines prepared therefrom which are able to trigger a protective immune response in...

...of cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...THE INVENTION

Applicants have discovered that cells of a pathogenic strain of a species of **Neospora** may be **attenuated**, and that the resulting attenuated cells are capable of triggering an immune response that protects...

...of cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...to any clinical symptom, condition, event or pathology associated with infection of the mammal by **Neospora**.

The term "**attenuated**" as used herein describes a cell, **culture**, or strain of **Neospora** exhibiting a detectable **reduction** in infectivity or virulence in vitro or in vivo as compared to that of the parent strain of **Neospora** from which the **attenuated** cell, culture, or strain is derived. Reduction in virulence encompasses any detectable decrease in any...

...or antigen that triggers a protective immune response when administered to a mammal.

Preparation Of **Attenuated** Strains Of **Neospora**

Since the invention is based on the discovery that cells of a pathogenic strain of **Neospora** may be **attenuated**, and that the resulting attenuated cells are capable of triggering an immune response that protects...

...exposure, is preferably that amount which results in producing one or more viable cells of **Neospora** that exhibit an **attenuated** level of pathogenicity but that are capable of triggering an immune response that protects against...

...for use of mutagenic agents may be determined empirically using standard techniques.

Pathogenic strains of **Neospora** may also be **attenuated** using

recombinant DNA technology according to techniques known in the art, and the present invention...

...not replicate sufficiently and will thus fail to produce plaques in host cell monolayers.

An **attenuated** strain of **Neospora** may be derived from any pathogenic strain of any species of the genus including, but...

...monkey kidney cells from the ATCC (ATCC Accession No. CRL-12230).

Both parental strains and **attenuated** strains of **Neospora** may be **cultured** in vitro by infecting any receptive cell line, preferably a mammalian cell line, with tachyzoites...

...according to known techniques described in the art. Mammalian cell lines in which tachyzoites of **Neospora** can be **cultured** include, for example, human foreskin fibroblasts (Lindsay et al., 1993, Am. J. Vet. Res. 54...

...Bradyzoites may be similarly cultured and manipulated.

Mammalian cell cultures can be grown, and cell **cultures** infected with **Neospora** can be maintained, in any one of several culture media described in the art. For...

...but in which the fetal bovine serum is increased to 10% (v/v) (growth medium). **Attenuated** strains of **Neospora** having novel auxotrophies will require appropriate modification to the culture medium to support growth, as known in the art.

Neospora -infected monolayer **cultures** of mammalian cells are typically maintained under standard tissue culture conditions such as, for example...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...and a veterinarily acceptable carrier.

The vaccine of the invention comprises live cells of an **attenuated** strain of **Neospora**, either as the sole antigenic component or in combination with one or more other antigens...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...formulations. Immunomodulatory agents are selected based on their ability to maintain both viability of the **attenuated Neospora** cells and ability of the cells to trigger a protective immune response in the vaccinated...

...term "protection" is used broadly and is not limited to absolute prevention of infection by **Neospora**, but includes a **reduction** in infectivity, or in the severity of a disease or condition resulting from infection, including...

...of NCTS-8 does not produce clinical disease.

The results indicate that a formulation comprising **attenuated**, live **Neospora** tachyzoites and an adjuvant is at least as effective and safe for use as a...

...the attenuated NCTS-8 strain.

Table 20 demonstrates the ability of a vaccine comprising live, **attenuated** tachyzoites of **Neospora** to protect pygmy goat does against

Neospora-induced abortion. All 4 goat does vaccinated with...

...SPECIFICATION invention relates to attenuated strains of the pathogenic protozoan, *Neospora*, and to live vaccines against **neosporosis** prepared from the **attenuated** strains which are useful in the prevention of clinical disease and abortion in mammals.

BACKGROUND...

...al., 1995, Am. J. Vet. Res. 56:1176-1180.

WO 9525541 discloses a biologically pure **culture** of bovine **Neospora**, methods of detecting anti-*Neospora* antibodies and *Neospora*-specific nucleic acids, and a composition containing...

...*Neospora* antigen and carrier for use as a vaccine. WO 9525541 does not, however, teach **attenuated** live **cultures** of **Neospora**, or live vaccines prepared therefrom which are able to trigger a protective immune response in...

...of cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...THE INVENTION

Applicants have discovered that cells of a pathogenic strain of a species of **Neospora** may be **attenuated**, and that the resulting attenuated cells are capable of triggering an immune response that protects...

...of cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...to any clinical symptom, condition, event or pathology associated with infection of the mammal by **Neospora**.

The term "**attenuated**" as used herein describes a cell, **culture**, or strain of **Neospora** exhibiting a detectable **reduction** in infectivity or virulence in vitro or in vivo as compared to that of the parent strain of **Neospora** from which the **attenuated** cell, culture, or strain is derived. Reduction in virulence encompasses any detectable decrease in any...

...or antigen that triggers a protective immune response when administered to a mammal.

Preparation Of **Attenuated** Strains Of **Neospora**

Since the invention is based on the discovery that cells of a pathogenic strain of **Neospora** may be **attenuated**, and that the resulting attenuated cells are capable of triggering an immune response that protects...

...exposure, is preferably that amount which results in producing one or

more viable cells of **Neospora** that exhibit an **attenuated** level of pathogenicity but that are capable of triggering an immune response that protects against...

...for use of mutagenic agents may be determined empirically using standard techniques.

Pathogenic strains of **Neospora** may also be **attenuated** using recombinant DNA technology according to techniques known in the art, and the present invention...

...not replicate sufficiently and will thus fail to produce plaques in host cell monolayers.

An **attenuated** strain of **Neospora** may be derived from any pathogenic strain of any species of the genus including, but...

...monkey kidney cells from the ATCC (ATCC Accession No CRL-12230).

Both parental strains and **attenuated** strains of **Neospora** may be **cultured** in vitro by infecting any receptive cell line, preferably a mammalian cell line, with tachyzoites...

...according to known techniques described in the art. Mammalian cell lines in which tachyzoites of **Neospora** can be **cultured** include, for example, human foreskin fibroblasts (Lindsay et al., 1993, Am. J. Vet. Res. 54...

...Bradyzoites may be similarly cultured and manipulated.

Mammalian cell cultures can be grown, and cell **cultures** infected with **Neospora** can be maintained, in any one of several culture media described in the art. For...

...but in which the fetal bovine serum is increased to 10% (v/v) (growth medium). **Attenuated** strains of **Neospora** having novel auxotrophies will require appropriate modification to the culture medium to support growth, as known in the art.

Neospora -infected monolayer **cultures** of mammalian cells are typically maintained under standard tissue culture conditions such as, for example...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...and a veterinarily acceptable carrier.

The vaccine of the invention comprises live cells of an **attenuated** strain of **Neospora**, either as the sole antigenic component or in combination with one or more other antigens...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora**, which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...formulations. Immunomodulatory agents are selected based on their ability to maintain both viability of the **attenuated Neospora** cells and ability of the cells to trigger a protective immune response in the vaccinated...

...term "protection" is used broadly and is not limited to absolute prevention of infection by **Neospora**, but includes a **reduction** in infectivity, or in the severity of a disease or condition resulting from infection, including...

...of NCT5-8 does not produce clinical disease.

The results indicate that a formulation comprising **attenuated**, live **Neospora** tachyzoites and an adjuvant is at least as effective and safe

for use as a...

...the attenuated NCTS-8 strain.

Table 20 demonstrates the ability of a vaccine comprising live, **attenuated** tachyzoites of **Neospora** to protect pygmy goat does against Neospora-induced abortion. All 4 goat does vaccinated with...

...CLAIMS of cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...CLAIMS of cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...live cells of a strain derived from a pathogenic parent strain of a species of **Neospora** , which cells exhibit **attenuated** pathogenicity compared to those of the parent strain but which are capable of triggering an...

...CLAIMS Neospora en vue de l'utilisation dans un vaccin qui protege un mammifere contre la **neosporose** , comprenant la **modification** des cellules provenant d'une souche parente pathogene d'une espece de Neospora; la selection...

...a 5.

12. Procede de preparation d'un vaccin pour proteger un mammifere contre la **neosporose** , comprenant la **modification** des cellules provenant d'une souche parente pathogene d'une espece de Neospora; la selection ...

2/3,KWIC/71 (Item 1 from file: 357)

DIALOG(R) File 357:Derwent Biotech Res.

(c) 2004 Thomson Derwent & ISI. All rts. reserv.

0318775 DBR Accession No.: 2003-19915 PATENT

New overexpressing homologous antigen vaccines, useful for immunization, prophylaxis or treatment of a vertebrate at risk of or suffering from, e.g. brucellosis, paratuberculosis, tuberculosis or neosporosis - vector expression in host cell for use in disease therapy and vaccine

AUTHOR: SCHURIG G; BOYLE S M; SRIRANGANATHAN N

PATENT ASSIGNEE: SCHURIG G; BOYLE S M; SRIRANGANATHAN N 2003

PATENT NUMBER: US 20030044431 PATENT DATE: 20030306 WPI ACCESSION NO.: 2003-521726 (200349)

PRIORITY APPLIC. NO.: US 268673 APPLIC. DATE: 20021011

NATIONAL APPLIC. NO.: US 268673 APPLIC. DATE: 20021011

LANGUAGE: English

...ABSTRACT: prophylaxis or treatment of a vertebrate at risk of or suffering from paratuberculosis, tuberculosis or **neosporosis** ,

comprises an **attenuated** or avirulent strain of a pathogenic bacteria of the genus *Brucella*, where the strain over...

2/3,KWIC/74 (Item 4 from file: 357)
DIALOG(R)File 357:Derwent Biotech Res.
(c) 2004 Thomson Derwent & ISI. All rts. reserv.

0238607 DBR Accession No.: 99-08708 PATENT
Vaccines containing *Neospora caninum* grown in tissue culture, useful for protecting cattle and dogs against infections and abortion - live attenuated subunit vaccine production
AUTHOR: Choromanski L; Brown K K
CORPORATE SOURCE: Pittsburgh, PA, USA.
PATENT ASSIGNEE: Bayer 1999
PATENT NUMBER: WO 9920303 PATENT DATE: 990429 WPI ACCESSION NO.: 99-288171 (9924)
PRIORITY APPLIC. NO.: US 954531 APPLIC. DATE: 971020
NATIONAL APPLIC. NO.: WO 98US21515 APPLIC. DATE: 981013
LANGUAGE: English

...ABSTRACT: *N. caninum*. (I) comprises an antigen selected from whole, inactivated or modified but live tissue **cultured *Neospora* spp.** tachyzoites or an extract or one or more subunits from these tachyzoites. (34pp)

DESCRIPTORS: ***Neospora caninum* tachyzoite live, attenuated**, subunit vaccine prep., tissue culture, appl. dog protection from abortion, infection mammal animal (Vol.18...

2/3,KWIC/75 (Item 5 from file: 357)
DIALOG(R)File 357:Derwent Biotech Res.
(c) 2004 Thomson Derwent & ISI. All rts. reserv.

0225574 DBR Accession No.: 98-07171 PATENT
Live attenuated *Neospora* vaccine - *Neospora caninum* culture for vaccine production
AUTHOR: Brake D A; Blagburn B L
CORPORATE SOURCE: New York, NY, USA; Auburn, AL, USA.
PATENT ASSIGNEE: Pfizer; Univ.Auburn 1998
PATENT NUMBER: NZ 329095 PATENT DATE: 980325 WPI ACCESSION NO.: 98-258985 (9823)
PRIORITY APPLIC. NO.: US 31248 APPLIC. DATE: 961112
NATIONAL APPLIC. NO.: NZ 97329095 APPLIC. DATE: 971103
LANGUAGE: English

Live attenuated *Neospora* vaccine - *Neospora caninum* culture for vaccine production

DESCRIPTORS: ***Neospora caninum* culture**, appl. **attenuated *Neospora* vaccine prep.**, dog, cow, goat, sheep, horse veterinary medicine protozoan mammal animal fermentation??? (Vol.17...

?logoff hold

16aug04 16:21:20 User228206 Session D2213.7
\$0.04 0.013 DialUnits File155
\$0.21 1 Type(s) in Format 9
\$0.21 1 Types
\$0.25 Estimated cost File155
\$0.07 0.013 DialUnits File5
\$3.50 2 Type(s) in Format 3
\$3.50 2 Types
\$3.57 Estimated cost File5
\$1.26 0.214 DialUnits File654
\$2.80 4 Type(s) in Format 3
\$2.80 4 Types
\$4.06 Estimated cost File654
\$0.46 0.097 DialUnits File349
\$8.00 5 Type(s) in Format 3

10507781 PMID: 10608449

Phenotypic characterisation of a *Neospora caninum* temperature-sensitive strain in normal and immunodeficient mice.

Dreier K J; Stewarter L W; Kerlin R L; Ritter D M; Brake D A

Animal Health Biological Discovery and Drug Safety Evaluation (RLK), Pfizer Central Research, Pfizer Inc., Groton, CT 06340, USA.

International journal for parasitology (ENGLAND) Oct 1999, 29 (10) p1627-34, ISSN 0020-7519 Journal Code: 0314024

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

The in vivo persistence, immunogenicity and pathogenicity of a recently described temperature-sensitive (ts) strain from ***Neospora caninum***, NCTs-8, was investigated in normal and immunodeficient mice. Groups of BALB/c and SCID/Bg mice were infected s.c. with 5×10^6 wild-type NC-1, control NCTs-8 (pass 0) or NCTs-8 tachyzoites prepared at four in vitro passage levels (pass 7, 13, 21 and 28). For persistence and immunogenicity studies, BALB/c mice were bled and sacrificed at 4, 6 or 8 weeks p.i. Sera were analysed by IFAT and brain tissues examined for lesions by histology and tested for parasite presence by PCR. For pathogenicity studies, SCID/Bg mice were monitored by clinical signs and survival time. Results from parasite persistence experiments demonstrated microscopic lesions and PCR positive brain tissues in NC-1 infected mice. In contrast, brain tissues from NCTs8-infected groups were consistently negative by histology and PCR. Based on IFAT titres, all parasite strains were immunogenic, although parasite-specific IgG levels were lower in the NCTs-8 infected groups. Results from pathogenicity studies in SCID/Bg mice demonstrated a significantly ($P < 0.0001$) longer mean survival time in NCTs-8 vs NC-1 infected groups. In addition, there was no significant difference in mean survival time between control NCTs-8 and experimental passage NCTs-8 infected mice. Collectively, these studies demonstrate that the NCTs-8 strain maintains a stable phenotype following multiple passages in vitro, and possesses an **attenuated**, shorter persistence phenotype in vivo compared with the parental wild-type NC-1.

Descriptors: Coccidiosis--parasitology--PS; * ***Neospora*** --physiology--PH; Animals; Antibodies, Protozoan--blood--BL; Brain--pathology--PA; DNA, Protozoan--analysis--AN; Mice; Mice, Inbred BALB C; Mice, SCID; ***Neospora*** --genetics--GE; ***Neospora*** --immunology--IM; ***Neospora*** --pathogenicity--PY; Phenotype; Polymerase Chain Reaction; Temperature; Virulence

CAS Registry No.: 0 (Antibodies, Protozoan); 0 (DNA, Protozoan)

Record Date Created: 20000120

Record Date Completed: 20000120

13431831 PMID: 9105320

Separation and cryopreservation of Neospora caninum tissue cysts from murine brain.

McGuire A M; McAllister M M; Jolley W R

Department of Veterinary Sciences, University of Wyoming, Laramie 82070, USA.

Journal of parasitology (UNITED STATES) Apr 1997, 83 (2) p319-21, ISSN 0022-3395 Journal Code: 7803124

Document type: Journal Article

Languages: ENGLISH

Main Citation Owner: NLM

Record type: Completed

Subfile: INDEX MEDICUS

A protocol was developed for the separation, concentration enumeration, and cryopreservation of **Neospora** caninum tissue cysts from mouse brains. Brains from chronically infected mice were homogenized and tissue cysts counted in 10-microliters aliquots. Tissue cysts were separated from brain homogenates by centrifugation at 4,400 g on 35% (v/v) Percoll/phosphate-buffered saline (PBS) continuous-density gradients. After removal of the brain layer, the separated tissue cysts were concentrated by diluting the remaining solution with PBS and centrifuging at 500 g. The pellet was resuspended in PBS and tissue cysts were enumerated. Fifty percent of tissue cysts were recovered from brains centrifuged once and 64% from brains centrifuged twice. Tissue cysts were preserved with 7.5% dimethyl sulfoxide in horse serum at -60 C. After thawing, bradyzoites were digested in an acid/pepsin solution and placed onto Vero cell cultures.

Neospora caninum tachyzoites were recovered from cell cultures, indicating that bradyzoites retained viability after concentration and cryopreservation. Separated tissue cysts ranged in diameter from 107 microns to 15 microns (average = 31 microns), and the average bradyzoite dimensions were 2 x 7.5 microns. These methods make it possible to store viable N. caninum tissue cysts for oral- **infectivity** trials and other studies.

Tags: Support, U.S. Gov't, Non-P.H.S.

Descriptors: Brain--parasitology--PS; *Cryopreservation; * **Neospora** --isolation and purification--IP; Animals; Centrifugation, Density Gradient ; Cercopithecus aethiops; Mice; Vero Cells

Record Date Created: 19970430